


# CORRELATION OF MAP UNITS



## DESCRIPTION OF MAP UNITS

#### Qal Alluvium, modern (Holocene)

Silty clay and sandy silt with minor sand and sparse gravel; thickness 10 to 30 feet (3 to 10 m); found along banks of Ohio River and in floodplains along streams tributary to the Ohio River; deposited by modern/historic stream processes; deposit is inset into adjacent map units; contact with adjacent units varies from sharp to poorly defined; mapped on the basis of topographic expression.

#### Qao

Alluvium, natural levee deposits (Holocene) Sand and silt; deposited in levee ridges or overwash deposits on floodplains of major rivers (Qafp) and on the Ohio River low outwash terraces (Qot1); grades into adjacent floodplain deposits; typically sandier than adjacent floodplain deposits.

Alluvium, active modern sloughs (Holocene) Organic-rich, black and gray clayey silt, silty clay, and clay; found within low lying areas on floodplain (Qafp) and low outwash terrace (Qot1); serve as poorly drained pathways to channel water off of the floodplain; areas that retain water year-round form bogs and cypress swamps.

## Qafp

Alluvium, Ohio River floodplain (Holocene) Sand, silt, fine gravel, and clay; surface mantled by silty clay and sandy silt; surface forms the lowest well-developed terrace along the Ohio River, 30 to 45 feet (10 to 15 m) thick; overlies sand and gravel deposits of older outwash deposits; contact is sharp, drawn at scarp of next higher terrace; estimated to range in age up to 6,500 years.

# Qafpg

Alluvium, Green River floodplain (Holocene) Silt, sand, fine gravel, and clay; surface mantled by silty clay and sandy silt; surface forms the lowest well-developed terrace along the Green River; 30 to 45 feet (10 to 15 m) thick; overlies older valley-fill units; contact is sharp, drawn at scarp of next higher terrace; estimated to range in age up to 6,500 years.

# Alluvium, modern slough, abandoned Green River meander (Holocene) Organic-rich, black and gray clayey silt, silty clay, and clay; deposited within recently abandoned

meander of Green River; can retain standing water for months; areas that retain water year-round form bogs and cypress swamps.

## Qafto

Alluvium, Ohio River floodplain terrace (Holocene) Sand, silt, fine gravel, and clay; surface mantled by silty clay and sandy silt; surface forms a distinct low relief terrace along the landward margin of the Ohio River floodplain (Qafp); 30 to 45 feet (10 to 15 m) thick; overlies sand and gravel deposits of older outwash deposits; contact is sharp, drawn at scarp of next higher terrace; estimated to range in age up to 6,500 years.

#### Qaftg

Alluvium, Green River floodplain terrace (Holocene) Silt, sand, fine gravel, and clay; surface mantled by silty clay and sandy silt; surface forms a distinct low relief terrace along the margin of the Green River floodplain (Qafpg); 30 to 45 feet (10 to 15 m) thick; overlies sand and gravel deposits of older outwash deposits; contact is sharp, drawn at scarp of next

#### higher terrace; estimated to range in age up to 6,500 years.

#### Qat1g Alluvium, low Green River terrace (Holocene)

Poorly sorted silt, sand, and clay deposited by the Green River; lithologically similar to Green River floodplain (Qafpg); distinguished by topographic expression from lower floodplain (Qafpg), but found below Ohio River low outwash terrace (Qot1).

Alluvium, young slough (Holocene), abandoned Green River channel (Pleistocene - Holocene) Clayey silt, silty sand, and silty clay; 30 to 45 feet (10 to 15 m) thick; forms arcuate, low-lying trough; represents an abandoned channel of Green River as it migrated across the low terrace (Qot1g); overlies older outwash deposits (Qot3); contact sharp, identified by surface topography; floods frequently.

#### Outwash, Ohio River scrollwork terrace (Pleistocene - Holocene) Qot1 o

Fine to coarse sand and gravel, with local lenses of silt and clay, gravel includes chert, quartzite, sandstone, siltstone, igneous and metamorphic rocks, limestone, and coal; lithologically similar to adjacent outwash terraces; surface mantled with alluvial silty sand and sandy silt; 30 to 45 feet (10 to 15 m) thick; surface forms well-developed, swell-and-swale topography on Ohio River low terrace; deposited as glacial outwash reworked during postglacial adjustment of the Ohio River; overlies older outwash deposits (Qot3); contact is approximate, inferred from surface topography.

# Alluvium, young slough (Holocene), abandoned Green River channel (Pleistocene - Holocene) Silty sand, clayey silt, and silty clay; 30 to 45 feet (10 to 15 m) thick; forms sinuous, low-lying trough (Katie Meadow Slough); represents an abandoned channel of Green River as it migrated across the low terrace (Qot1g); overlies older outwash deposits (Qot3); contact sharp, identified by surface topography; floods frequently.

#### Qot1g

Outwash, Green River scrollwork terrace (Pleistocene - Holocene) Fine to coarse sand and gravel, with local lenses of silt and clay, gravel includes chert, quartzite, sandstone, siltstone, limestone, and coal; lithologically similar to adjacent outwash terraces; surface mantled with alluvial silty sand and sandy silt; 30 to 45 feet (10 to 15 m) thick; surface forms well-developed, swell-and-swale topography on Ohio River low terrace; deposited as point bar deposits of glacial outwash reworked by meandering postglacial Green River; overlies older outwash deposits (Qot3); contact is approximate, inferred from surface topography.

#### Qot1

Outwash, low terrace (Pleistocene - Holocene) Fine to coarse sand and gravel, with local lenses of silt and clay; gravel includes chert, quartzite, sandstone, siltstone, igneous and metamorphic rocks, limestone, and coal; lithologically similar to adjacent outwash terraces (Qot2, Qot3); surface mantled with alluvial silty sand and sandy silt; 30 to 45 feet (10 to 15 m) thick; surface forms well-developed, low-relief terrace along Ohio River valley; deposited as glacial outwash reworked by post-glacial Ohio River; overlies older outwash deposits (Qot3); contact is sharp, drawn at scarp of next higher terrace or upland.

## Qel

Loess (Pleistocene-Holocene) (thicker than ~3 to 5 feet) Silt, clayey silt, and fine sand deposited by wind; typically massive; unit thickest (up to 40 feet) near Ohio River valley and thins gradually to the south; mantles bedrock upland; not mapped where locally found on lacustrine terrace (Qlt) and high outwash terraces (Qot3); estimated to range in age from 22,500 to 10,000 years old; locally includes thin layers of loess inferred to be older than 30,000 years.

Alluvium, slough (Holocene), abandoned Green River channel (Pleistocene) Clayey silt, silty clay, and silty sand; 30 to 45 feet (10 to 15 m) thick; forms sinuous, low-lying trough inset into Green River paleovalley (Qapg); represents an abandoned channel of Green River as it migrated across the high terrace (Qot3); overlies older outwash (Qot3); contact sharp, identified by surface topography; floods occasionally.

#### Qapg

Green River paleovalley (Pleistocene) Silty sand, clayey silt and silty clay with minor chert gravel; 30 to 45 feet (10 to 15 m) thick; includes Beds at Hubert Court of Ray (1965); forms broad, linear trough inset into and overlying deposits of adjacent high outwash terrace (Qot3) and lacustrine terrace (Qlt); represents abandoned Pleistocene paleovalley of the Green River; contact is sharp, drawn at scarp of adjacent high outwash or lacustrine terrace; wood from about 40 feet deep has been radiocarbon dated to 23,150 ± 500 ypb (Ray, 1965).

#### Qlt Lacustrine terrace (Pleistocene)

Clayey silt and silty clay; 30 to 45 feet (10 to 15 m) thick, thicker in tributary valleys; overlying complex deposits of sand, silt, clay and minor gravel; locally mantled by loess (Qel, not mapped); forms prominent low-relief terrace in tributary valleys and sheltered portions of Ohio River valley; unit deposited in lacustrine and slackwater environments associated with alluviation of the Ohio River valley by glacial outwash and resulting impoundment of tributary valleys; underlying material is of apparent mixed fluvial and fluvio-lacustrine origin; contact with fluvial units is sharp, and drawn on scarps separating adjacent terraces; contact with eolian and upland units (Qel, Qes) is gradational and approximate, inferred by surface topography; estimated to rage in age from 23,000 to 18,000 years old.

#### Qltm

Marginal lacustrine deposits (Pleistocene) Clayey silt, silt, and fine sand; thickness uncertain; surface forms moderate slope and benched upland areas bordering lacustrine deposits (Qlt); represents complex transition between lacustrine deposits and loess mantling upland; deposits include loess, loess-derived slopewash, colluvium, lacustrine silt and clay, and lacustrine shoreline deposits; contacts gradational and approximate, mapped on the basis of topographic expression.

#### Qot3

Outwash, high terrace (Pleistocene)) Fine to coarse sand and gravel, with local lenses of silt and clay; gravel includes chert, quartzite, sandstone, siltstone, igneous and metamorphic rocks, limestone, and coal; lithologically similar to adjacent outwash terraces; surface mantled with eolian and alluvial silty sand and sandy silt; up to 170 feet (52 m) thick; surface forms well-developed, dissected terrace along Ohio River valley; deposited as glacial outwash; represents maximum valley filling by glacial outwash valley train deposits; overlies bedrock (Pz) or older pre-glacial alluvial deposits (not snown); contact is sharp, drawn at scarp of adjacent terrace or upland; age estimated to be 120,000 to 22,000 years old.

af3

Upland gravel (Pliocene-Pleistocene) Gravel and medium to coarse sand; pebbles include brown, patina chert, quartz, and silicified fossils; locally cemented by iron oxide; thickness uncertain; unit found on uplands, covered by loess and poorly exposed; comparable to the Luce Gravel of Ray (1965).

Pz Bedrock (Paleozoic) Consolidated shale, sandstone, coal, and overlying poorly sorted regolith, comprising the core of the uplands in the study area; includes areas of loess thinner than 3 ft (1 m).

Artificial fill, engineered fill (Modern) af1 Unconsolidated material used as fill for the construction of roads, railroads, buildings, and floodwalls.

Artificial fill, mine spoil (Modern) Disturbed bedrock and regolith produced from mining operations. af2

> Artificial fill, other (Modern) Chaotic, unconsolidated fill material; includes material dredged from creeks to form artificial levees.

New water (Modern) nw Areas of former land which have been removed by active erosion or dredging since the completion of topographic mapping.

#### EXPLANATION

State boundary	5 1
River bank	Roads NV Parkway or Federal highway
Contact, observed	State roads
Contact, approximately located	Local roads
Contact, inferred	Local roads - gravel
Contact (concealed)	angeo de mensiona entre
Faults (concealed)	Surface observations ▲ Landform only
Scarp, teeth point downward	Lithologic data
Railroads	<ul> <li>KGS databases</li> <li>Cross section location</li> </ul>
Mined areas	
	River bank Contact, observed Contact, approximately located Contact, inferred Contact (concealed) Faults (concealed) Scarp, teeth point downward Railroads

#### ACKNOWLEDGEMENTS

This map was generated using new field mapping and compilation of unpublished and previously published data and was funded in part by the U.S. Geological Survey National Cooperative Geologic Mapping Program under the STATEMAP Program authorized by the National Geologic Mapping Act of 1992, Grant No. 05HQAG0007, and by the Kentucky Geological Survey.

Field mapping was completed by Mark F. Thompson and Michael M. Murphy from April 2005 to March 2006, with assistance from William M. Andrews Jr., Ronald C. Counts and Steven L. Martin (KGS).

Subsurface information was compiled from data on file at the Kentucky Geological Survey.

## REFERENCES

Cox, Frank R., Jr.,1974, Soil survey Daviess and Hancock Counties, Kentucky; U.S. Department of Agriculture, 108p.

Converse Henry T., Jr., and Cox, Frank R., Jr., 1967, Soil survey of Henderson County, Kentucky; U.S. Department of Agriculture, 108p.

Johnson, W.D., Jr., 1972, Geologic map of the Newburgh and Yankeetown quadrangles, Henderson and Daviess Counties, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1045, scale 1:24,000.

Johnson, W.D., Jr., 1972, Geologic map of the Reed quadrangle, Kentucky-Indiana: U.S. Geological Survey Geologic Quadrangle Map GQ-1038, scale 1:24,000.

Ray, L.L., 1965, Geomorphology and Quaternary geology of the Owensboro quadrangle, Indiana and Kentucky: U.S. Geological Survey Professional Paper 488, 71p.

Solis, M.P., and Terry, J., 2000, Spatial database of the Newburgh and Yankeetown quadrangles, Henderson and Daviess Counties, Kentucky: Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-1045. Adapted from Johnson, W.D., Jr., 1972, Geologic map of the Newburgh and Yankeetown quadrangles, Henderson and Daviess Counties, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1045, scale 1:24,000.

QUADRANGLE INDEX


Solis, M.P., and Terry, J., 2000, Spatial database of the Reed quadrangle, Kentucky-Indiana: Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-1038. Adapted from Johnson, W.D., Jr., 1972, Geologic map of the Reed quadrangle, Kentucky-Indiana: U.S. Geological Survey Geologic Quadrangle Map GQ-1038, scale 1:24,000.

Theis, Charles Vernon, 1929, The Geology of Henderson County, Kentucky, University of Cincinnati, unpublished dissertation, 251p.

DISCLAIMER

Although these data have been processed successfully on a computer system at the Kentucky Geological Survey (KGS), no warranty, expressed or implied, is made by KGS regarding the utility of the data on any other system, nor shall the act of distribution constitute any such warranty. KGS does not guarantee this map or digital data to be free of errors or inaccuracies. KGS disclaims any responsibility or liability for interpretations from this map or digital data, or decisions based thereon.

The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Government.



QUATERNARY GEOLOGIC MAP OF PARTS OF THE YANKEETOWN AND REED 7.5-MINUTE QUADRANGLES, WESTERN KENTUCKY

By Mark F. Thompson, Michael L. Murphy, and William M. Andrews Jr.