

CORRELATION OF MAP UNITS

DESCRIPTION OF MAP UNITS

- Qaf** Alluvium, modern (Holocene)
Silty clay and sandy silt with minor sand and sparse gravel; thickness 10 to 30 feet (3 to 10 m); found along river banks and in floodplains of smaller streams; deposited by modern/historic stream processes; deposit is inset into adjacent map units; contact with adjacent units varies from sharp to poorly defined; locally inferred on the basis of topographic expression. Some streams in the mapped area have been rerouted for land use purposes; locally, some Qaf dredged from these streams has been extensively redistributed across adjacent fields and is unmappable.
- Qao** Alluvium, natural levee deposits (Holocene)
Sand and silt; deposited by levees or overwash deposits on floodplains of major rivers (Qaf) and on the Ohio River low-lying terraces (Qot1); grades into adjacent floodplain deposits; typically sandier than adjacent floodplain deposits.
- Qas** Alluvium, active modern floodplain sloughs (Holocene)
Organic-rich, black and gray clayey silt, silty clay, and clay; found within low lying areas on floodplain (Qaf) and low-lying terraces (Qot1); serve as poorly drained pathways which channel water from the floodplain; areas that retain water year-round form bogs and cypress swamps.
- Qat** Alluvium, alluvial fans (Holocene)
Silt, sand, and gravel; thickness uncertain; forms fan-shaped alluvial-colluvial aprons at mouths of small valleys; deposited by floods and debris flows from small tributary valleys developed in loess-mantled uplands; extent of unit mapped by topographic expression.
- Qoc** Colluvium (Holocene)
Silt, sand, clay, and rock fragments; unsorted; which has been transported downslope under the influence of gravity; primarily mantles steep slopes.
- Qatp** Alluvium, river floodplains (Holocene)
Silt, sand, fine gravel, and clay; surface mantled by silty clay and sandy silt; surface forms the lowest well-developed terrace along major rivers; 30 to 45 feet (10 to 15 m) thick; overlies older unconsolidated deposits or bedrock; contact is sharp, drawn at scarp of next higher terrace; estimated to range in age up to 6,500 years.
- Qas1** Alluvium, abandoned Green River meander (Holocene)
Organic-rich, black and gray clayey silt, silty clay, and clay; deposited within recently abandoned meander of Green River; can retain standing water for months; areas that retain water year-round form bogs and cypress swamps.
- Qat** Alluvium, low terrace (Holocene)
Silt, sand, and clay deposited by rivers; forms terrace above adjacent floodplain (Qaf); contact with adjacent units varies from sharp to poorly defined; locally inferred on the basis of topographic expression; distinguished by topographic expression from lower floodplain (Qaf) but found below Ohio River low-lying terrace (Qot1) and lacustrine terrace (Qit).
- Qas2** Alluvium, abandoned Green River channel (Pleistocene - Holocene)
Clayey silt, silty sand, and silty clay; 30 to 45 feet (10 to 15 m) thick; forms arcuate, low-lying trough; represents an abandoned channel of Green River as it migrated across the low terrace (Qot1g); overlies older outwash deposits (Qot2); contact sharp, identified by surface topography; floods frequently.
- Qot1** Alluvium, reworked outwash, Ohio River scrollwork terrace (Pleistocene - Holocene)
Fine to coarse sand and gravel, with local lenses of silt and clay; gravel includes chert, quartzite, sandstone, siltstone, igneous and metamorphic rocks, limestone, and coal; lithologically similar to adjacent outwash terraces; surface mantled with alluvial silty sand and sandy silt; 30 to 45 feet (10 to 15 m) thick; surface forms well-developed, well-sorted topography on Ohio River low terrace; worked during postglacial adjustment of the Ohio River; overlies older outwash deposits (Qot2); contact is approximate, inferred from surface topography.
- Qas3** Alluvium, abandoned Green River channel (Pleistocene - Holocene)
Silty sand, clayey silt, and silty clay; 30 to 45 feet (10 to 15 m) thick; forms sinuous, low-lying trough (Katie Meadow Slough); represents an abandoned channel of Green River as it migrated across the low terrace (Qot1g); overlies older outwash deposits (Qot2); contact sharp, identified by surface topography; floods frequently.

EXPLANATION

- Contact
- Approximate contact
- Fault
- Inferred fault

GEOLOGIC SUMMARY

GEOLOGIC SETTING

The regional project area is located in the lower Ohio River Valley, and includes the confluence of the Green River with the Ohio River. The landscape of the map area is characterized by low-relief bedrock uplands separated by broad alluvial valleys. Although the area is south of the Pleistocene glacial limit, the Ohio River served as a major outlet for glacial meltwater and entrained sediment during glacial stages. Rapid accumulation of glacial outwash in the main Ohio River Valley and along the mouths of tributaries led to impoundment and extensive deposition of slackwater and lacustrine sediment in the alluvial valleys. This lacustrine deposit has a complex and gradational transition with loess mantling adjacent uplands. The loess was primarily derived from the valley-bottom outwash. The uplands are underlain by Pennsylvanian relatively flat-lying, coal-bearing strata.

GEOTECHNICAL BEHAVIOR

The Quaternary deposits identified in the map area exhibit a wide range of grain size and geotechnical behaviors. Grain size distribution is one of the primary factors affecting the behavior of soils for geotechnical, hydrogeological, and agricultural applications. The grain size distribution of unconsolidated sediments is dominantly controlled by the conditions under which the material was deposited. Low energy environments allow the deposition of fine-grained materials. High energy deposits limit deposition to only coarser grained materials. Eolian processes produce very well sorted (poorly graded) materials. Fluvial processes produce moderate sorting; colluvial processes produce poorly sorted deposits.

HAZARDS

Flooding is a nearly annual occurrence along the Ohio River. Floods in the late winter or early spring commonly inundate low-lying areas in the floodplain. Larger floods occur roughly every 10 to 20 years (e.g. 1913, 1945, 1964, 1997), and cover parts of the low terraces. The maximum flood of record in the valley was in 1937, flooding river towns throughout the valley. Only structures on the highest outwash terraces and the lacustrine terrace (Qit) were spared flood damage. The impact of flooding is reflected in land-use patterns through the area. Older homes and businesses have survived on the lacustrine and high outwash terraces, and on the highest parts of low terraces (Qot1, Qot1g, Qot1g). Trailers and less expensive built homes are constructed on the low terraces. Only barns are found on the high parts of the floodplains (Qaf). The floodplain and low parts of the low terraces are dominantly left to woodlands or used for row-crop agriculture. Most livestock husbandry in the alluvial valleys has been abandoned and is now restricted to upland areas above the 10- to 20-year flood zone. The low-relief lacustrine terrace is locally very poorly drained.

The silt soils that dominate the loess-mantled uplands are highly erodible. Soil piping and associated cover collapses are common hazards as ground water seeps through the silt and is commonly perched above fragrans. Great care must be taken during agricultural operations not to mobilize and lose this valuable resource.

The map area is proximal to the Wabash Valley Seismic Zone and the New Madrid Seismic Zone. Small earthquakes have been felt in the area relatively frequently. The significant thickness of unconsolidated sediment (locally as much as 140 feet in the regional map area) raise concerns about ground motion amplification of seismic waves and potential liquefaction. The variations in lithology and thickness between materials in different map units will likely cause different responses of these materials to seismic shaking.

Qot1g Alluvium, reworked outwash, Green River scrollwork terrace (Pleistocene - Holocene)
Fine to coarse sand and gravel, with local lenses of silt and clay; gravel includes chert, quartzite, sandstone, siltstone, igneous and metamorphic rocks, limestone, and coal; lithologically similar to adjacent outwash terraces; surface mantled with alluvial silty sand and sandy silt; 30 to 45 feet (10 to 15 m) thick; surface forms well-developed, well-sorted topography on Ohio River low terrace; deposited as point bar deposits by meandering postglacial Green River; overlies older outwash deposits (Qot2); contact is approximate, inferred from surface topography.

Qot1 Alluvium, outwash, low terrace (Pleistocene - Holocene)
Silt, sand, and clay deposited by rivers; forms terrace above adjacent floodplain (Qaf); contact with adjacent units varies from sharp to poorly defined; locally inferred on the basis of topographic expression; distinguished by topographic expression from lower floodplain (Qaf) but found below Ohio River low-lying terrace (Qot1) and lacustrine terrace (Qit).

Qas2 Alluvium, abandoned Green River channel (Pleistocene)
Clayey silt, silty sand, and silty clay; 30 to 45 feet (10 to 15 m) thick; forms sinuous, low-lying trough (Katie Meadow Slough); represents an abandoned channel of Green River as it migrated across the low terrace (Qot1g); overlies older outwash deposits (Qot2); contact is sharp, drawn at scarp of next higher terrace or upland; floods occasionally.

Qas3 Alluvium, abandoned Green River channel (Pleistocene)
Clayey silt, silty clay, and silty sand; 30 to 45 feet (10 to 15 m) thick; forms sinuous, low-lying trough inset into Green River paleovalley (Qap); represents an abandoned channel of Green River as it migrated across the high terrace (Qot2); overlies older outwash deposits (Qot2); contact sharp, identified by surface topography; floods occasionally.

Qap Alluvium, Green River paleovalley (Pleistocene)
Silty sand, clayey silt and silty clay with minor chert gravel; 30 to 45 feet (10 to 15 m) thick; includes Beck at Hutter Court of Ray (1965); forms broad, linear trough inset into and overlying deposits of adjacent high outwash terrace (Qot2) and lacustrine terrace (Qit); represents abandoned Pleistocene paleovalley of the Green River; contact is sharp, drawn at scarp of adjacent high outwash or lacustrine terrace; wood from about 40 feet deep has been radiocarbon dated to 23,150 ± 500 ypb (Ray, 1965).

Qitn Upland marginal lacustrine deposits (Pleistocene)
Clayey silt, silt, and fine sand; thickness uncertain; surface forms moderate slope and bench upland areas bordering lacustrine deposits (Qit); represents complex transition between lacustrine deposits and loess mantling upland; deposits include loess, loess-derived slopewash, colluvium, lacustrine silt and clay, and lacustrine shoreline deposits; contacts gradational and approximate; mapped on the basis of topographic expression.

Qit Slackwater deposits, lacustrine terrace (Pleistocene)
Clayey silt and silty clay; 30 to 45 feet (10 to 15 m) thick; thicker in tributary valleys; overlying complex deposits of sand, silt, clay and minor gravel; locally mantled by loess (similar to Qot, not mapped); forms prominent low-relief terrace in tributary valleys and sheltered portions of Ohio River valley; unit deposited in lacustrine and slackwater environments associated with alluviation of the Ohio River valley by glacial outwash and resulting impoundment of tributary valleys; underlying material is of apparent mixed fluvial and fluvio-lacustrine origin; contact with fluvial units is sharp, drawn on scarp separating adjacent terraces; contact with colluvium and upland units (Qot, Qes, Qitn) is gradational and approximate, inferred by surface topography; estimated to range in age from 23,000 to 18,000 years old.

Qitg Upland gravel (Pleistocene-Pleistocene)
Gravel and medium to coarse sand; pebbles include brown, patina chert, quartz, and silicified fossils; locally cemented by iron oxide; thickness uncertain; unit found on uplands, covered by loess and poorly exposed; comparable to the Lacey Gravel of Ray (1965).

Pz Bedrock and residuum (Paleozoic)
(cross-section only) Consolidated shale, sandstone, coal, and overlying poorly sorted regolith, comprising the core of the uplands in the study area; includes areas of loess thinner than 3 to 5 ft (1 to 1.6 m).




af1 Artificial fill, engineered fill (Modern)
Compacted material used as fill for the construction of roads, railroads, buildings, floodwalls, and other engineered structures. Present in all areas of development; mapped only where fill distinctly changes the elevation.

af2 Artificial fill, mine spoil (Modern)
Disturbed bedrock and regolith produced from mining operations.

af3 Artificial fill, other (Modern)
Chaotic, unconsolidated fill material; includes material dredged from creeks to form artificial levees. Mapped only where fill is distinct.

nw New water (Modern)
Areas of former land which have been removed by active erosion or dredging since the completion of original topographic mapping.

Universal Transverse Mercator projection, zone 16, North American Datum of 1927
Topographic base and cultural features are Kentucky Raster Graphics (KRG) from Kymartian.ky.gov/kgmaps/KRG of Mount Vernon



**QUATERNARY GEOLOGIC MAP OF PART OF THE MOUNT VERNON
7.5-MINUTE QUADRANGLE, WESTERN KENTUCKY**
By Ronald C. Counts
2007