



88°00'00" 37°45'00"

**CORRELATION OF MAP UNITS**

Qaf	Qc	Qai	Qao	Qas	Qa
Qafp	Qafq	Qaf1	Qaf2	Qaf3	Qaf4
Qaf					
Qel	Qes	Qet	Qeo	Qeg	Qem
Qel	Qet	Qeo	Qem		
Qel	Qes	Qet	Qeo	Qeg	Qem
Qel	Qes	Qet	Qeo	Qeg	Qem

**DESCRIPTION OF MAP UNITS**

**Qai Alluvium, modern (Holocene)**  
Silty clay and sandy silt with minor sand and sparse gravel; thickness 10 to 30 feet (3 to 10 m); found along river banks and in floodplains of smaller streams; deposited by modern/historic stream processes; deposit is inset into adjacent map units; contact with adjacent units varies from sharp to poorly defined; locally inferred on the basis of topographic expression. Some streams in the mapped area have been routed for land-use purposes; locally, some Qai dredged from these streams has been extensively redistributed across adjacent fields and is unmappable.

**Qao Alluvium, natural levee deposits (Holocene)**  
Sand and silty sand deposited in levee ridges or overwash deposits on floodplains of major rivers (Qafp) and on the Ohio River low-terrace (Qa1); grades into adjacent floodplain deposits; typically sandier than adjacent floodplain deposits.

**Qas Alluvium, active modern floodplain sloughs (Holocene)**  
Organic-rich, black and gray clayey silt, silty clay, and clay; found within low-lying areas on floodplain (Qafp) and low-terrace (Qa1); serve as poorly drained pathways which channel water from the floodplain; areas that retain water year-round form bogs and cypress swamps.

**Qaf Alluvium, alluvial fans (Holocene)**  
Silt, sand, and gravel; thickness uncertain; forms fan-shaped alluvial-coluvial aprons at mouths of small valleys; deposited by floods and debris flows from small tributary valleys developed in loess-mantled uplands; extent of unit mapped by topographic expression.

**Qc Colluvium (Holocene)**  
Silt, sand, clay, and rock fragments; unsorted; which has been transported downslope under the influence of gravity; primarily mantles steep slopes.

**Qafp Alluvium, river floodplains (Holocene)**  
Sand, silt, fine gravel, and clay; surface mantled by silty clay and sandy silt; surface forms the lowest well-developed terrace along major rivers; 30 to 45 feet (10 to 15 m) thick; overlies older unconsolidated deposits or bedrock; contact is sharp, drawn at scarp of next higher terrace; estimated to range in age up to 6,500 years.

**Qa1 Alluvium, abandoned Green River meander (Holocene)**  
Organic-rich, black and gray clayey silt, silty clay, and clay; deposited within recently abandoned meander of Green River, can retain standing water for months; areas that retain water year-round form bogs and cypress swamps.

**Qafq Alluvium, low terrace (Holocene)**  
Silt, sand, and clay deposited by rivers; forms terrace above adjacent floodplain (Qafp); contact with adjacent units varies from sharp to poorly defined; locally inferred on the basis of topographic expression; distinguished by topographic expression from lower floodplain (Qafp), but found below Ohio River low-terrace (Qa1) and lacustrine terrace (Ql).

**Qa2 Alluvium, abandoned Green River channel (Pleistocene - Holocene)**  
Clayey silt, silty sand, and silty clay; 30 to 45 feet (10 to 15 m) thick; forms arcuate, low-lying trough (Katie Meadows Slough); represents an abandoned channel of Green River as it migrated across the low terrace (Qafq); overlies older outwash deposits (Qa2); contact sharp, identified by surface topography; floods frequently.

**Qa3 Alluvium, reworked outwash, Ohio River scrowork terrace (Pleistocene - Holocene)**  
Fine to coarse sand and gravel, with local lenses of silt and clay; gravel includes chert, quartzite, sandstone, siltstone, igneous and metamorphic rocks, limestone, and coal; lithologically similar to adjacent outwash terraces; surface mantled with alluvial silty sand and sandy silt; 30 to 45 feet (10 to 15 m) thick; surface forms well-developed, well-and-swale topography on Ohio River low terrace; reworked during postglacial adjustment of the Ohio River; overlies older outwash deposits (Qa2); contact is approximate, inferred from surface topography.

**Qa4 Alluvium, abandoned Green River channel (Pleistocene - Holocene)**  
Silty sand, clayey silt, and silty clay; 30 to 45 feet (10 to 15 m) thick; forms sinuous, low-lying trough (Katie Meadows Slough); represents an abandoned channel of Green River as it migrated across the low terrace (Qafq); overlies older outwash deposits (Qa2); contact sharp, identified by surface topography; floods frequently.

**Qa5 Alluvium, reworked outwash, Green River scrowork terrace (Pleistocene - Holocene)**  
Fine to coarse sand and gravel, with local lenses of silt and clay; gravel includes chert, quartzite, sandstone, siltstone, igneous and metamorphic rocks, limestone, and coal; lithologically similar to adjacent outwash terraces; surface mantled with alluvial silty sand and sandy silt; 30 to 45 feet (10 to 15 m) thick; surface forms well-developed, well-and-swale topography on Ohio River low terrace; deposited as point bar deposits by meandering postglacial Green River; overlies older outwash deposits (Qa2); contact is approximate, inferred from surface topography.

**GEOLOGIC SUMMARY**

**GEOLOGIC SETTING**

The regional project area is located in the lower Ohio River Valley, downstream of the confluence of the Wabash River and Ohio River. The landscape of the map area is characterized by very low to high-relief bedrock uplands separated by broad valleys. Although the area is just south of the Pleistocene (Illinoian) glacial limit, both the Ohio and Wabash Rivers served as major outlets for glacial meltwater and entrained sediment during glacial stages. Rapid accumulation of glacial outwash in the valleys and along the mouths of tributaries led to impoundment and extensive deposition of slackwater and lacustrine sediment in the tributary valleys. This lacustrine deposit has a complex and gradational transition with loess mantling adjacent uplands. The loess was primarily derived from the valley-bottom outwash. The uplands are underlain by faulted Upper Mississippian limestone and Pennsylvanian coal-bearing strata steeply dipping North to Northeast.

**GEOTECHNICAL BEHAVIOR**

The Quaternary deposits identified in the map area exhibit a wide range of grain size and geotechnical behaviors. Grain size distribution is one of the primary factors affecting the behavior of soils for geotechnical, hydrogeologic, and agricultural applications. The grain size distribution of unconsolidated sediments is dominantly controlled by the conditions under which the material was deposited. Low energy environments allow the deposition of fine-grained materials. High energy deposits limit deposition to only coarser grained materials. Eolian processes produce very well sorted (poorly graded) materials. Fluvial processes produce moderate sorting; colluvial processes produce poorly sorted deposits.

**HAZARDS**

Flooding is a nearly annual occurrence along the Ohio River. Floods in the late winter or early spring commonly inundate low-lying areas in the floodplain. Larger floods occur roughly every 10 to 20 years (e.g., 1913, 1945, 1964, 1997, 2007), and cover parts of the low terraces. The maximum flood of record in the Ohio River valley was in 1937; flooding river towns throughout the valley. Only structures on the highest outwash terraces and the lacustrine terrace (Ql) were spared flood damage. The impact of flooding is reflected in land-use patterns through the area. Older homes and businesses have survived on the lacustrine and high outwash terraces, and on the highest parts of low terraces (Qa1, Qa10, Qa11). Trailers and less expensive built homes are constructed on the low terraces. Only barns are found on the high parts of the floodplain (Qafp). The floodplain and low parts of the low terraces are dominantly left to woodlands or used for low-emp agriculture. Most livestock husbandry in the alluvial valleys has been abandoned and is now restricted to upland areas above the 10- to 20-year flood zone. The low-relief lacustrine terrace is locally very poorly drained.

The salt soils that dominate the loess-mantled uplands are highly erodible. Soil piling and associated cover collapses are common hazards as ground water seeps through the soil and is commonly perched above fragipans. Great care must be taken during agricultural operations not to mobilize and lose this valuable resource.

The map area is proximal to the Wabash Valley Seismic Zone, the New Madrid Seismic Zone, and is within the Rough Creek - Shawneetown Fault Zone. Small to moderate earthquakes have been felt in the area relatively frequently. The significant thickness of unconsolidated sediment (locally as much as 140 feet in the regional map area) raise concerns about ground motion amplification of seismic waves and potential liquefaction. The variations in lithology and thickness between materials in different map units will likely cause different responses of these materials to seismic shaking.

**EXPLANATION**

**Q1** KGS database, number indicate depth to bedrock in feet  
**Q2** Inferred Contact  
**Q3** KGS drilling, number indicate depth to bedrock in feet  
**Q4** KYTC data, number indicate depth to bedrock in feet  
**Q5** Landform observation and soil probe  
**Q6** Landform observation

**DISCLAIMER**

Although these data have been processed successfully on a computer system at the Kentucky Geological Survey (KGS), no warranty, expressed or implied, is made by KGS regarding the utility of the data on any other system, nor shall the act of distribution constitute any such warranty.

KGS does not guarantee this map or digital data to be free of errors or inaccuracies. Some cultural features originate from data sources other than KGS, and may not align with geologic features on this map. KGS disclaims any responsibility or liability for interpretations from this map or digital data, or decisions based thereon.

The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Government.

**ACKNOWLEDGMENTS**

This map was generated using new field mapping and compilation of unpublished and previously published data and was funded in part by the U.S. Geological Survey National Cooperative Mapping Program under the STATEMAP Program authorized by the National Geographic Mapping Act of 1992, Grant No. 08HQPM003, and by the Kentucky Geological Survey. Field mapping was completed by Scott Waninger from October 2008 to June 2009, with assistance from R. Counts (KGS).

Subsurface information was compiled from data on file at the Kentucky Geological Survey as well as data contributed by the Kentucky Transportation Cabinet and the U.S. Geological Survey.

**REFERENCES**

Crawford, M.M., 2004. Spatial database of the Grove Center quadrangle, Kentucky-Illinois, and part of the Shawneetown quadrangle, Kentucky. Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVQ-GQ-1314. Adapted from Palmer, J.E., 1976. Geologic map of the Grove Center quadrangle, Kentucky-Illinois, and part of the Shawneetown quadrangle, Kentucky-Illinois. U.S. Geological Survey Geologic Quadrangle Map GQ-1314, scale 1:24,000.

Faust, R.J., Barnfield, G.R., and Willinger, G.A., 1980. A Compilation of Ground Water Quality Data for Kentucky. U.S. Geological Survey, Open File Report 80-685, p. 963.

Frye, J.C., Leonard, A.B., Willman, H.B., and Glans, H.D., 1972. Geology and Paleontology of Late Pleistocene Lake Saline, Southeastern Illinois; Illinois State Geological Survey, Circular 471, 44p.

Glenn, L.C., 1912. Coals of the Tradewater River Region, Kentucky Geological Survey, Bulletin #17, Serial #24, 75p.

Glenn, L.C., 1922. The Geology and Coals of Webster County, Kentucky Geological Survey, Series #6, Vol #5, 249p.

Heinrich, P.V., 1982. Geomorphology and Sedimentology of Pleistocene Lake Saline, Southern Illinois. M.S. Thesis, University of Illinois, Urbana-Champaign, 145p.

Jacobs, E.H., 1981. Soil survey of Union and Webster Counties, Kentucky. U.S. Department of Agriculture, p. 126.

Jilson, W.R., 1943. The Geology of Union County, Kentucky, The Standard Printing Company, 83p.

McFarlan, A.C., 1961. Geology of Kentucky; University of Kentucky; Kentucky Geological Survey, 53p.

Mull, D.S., Cushman, R.V., and Lambert, T.W., 1971. Public and Industrial Water Supplies of Kentucky 1968-69. Kentucky Geological Survey Series 10, IC 20 p.107.

Nelson, W.J. and Lamm, D.K., 1986. Geologic Map of the Shawneetown Quadrangle, Gallatin County Illinois. Illinois Geological Survey, Map 104-1.

Palmer, J.E., 1976. Geologic map of the Grove Center quadrangle, Kentucky-Illinois, and part of the Shawneetown quadrangle, Kentucky. U.S. Geological Survey Geologic Quadrangle Map GQ-1314, scale 1:24,000.

Ray, L.L., 1965. Geomorphology and Quaternary geology of the Owensboro quadrangle, Indiana and Kentucky. U.S. Geological Survey Professional Paper 488, 71p.

Theis, C.V., 1929. The Geology of Henderson County, Kentucky, University of Cincinnati, 251p.

