

GEOLOGIC SETTING

The regional project area is located in the lower Ohio River valley, downstream of the confluence of the Wabash River and Ohio River. The landscape of the map area is characterized by very low to high-relief bedrock uplands separated by broad valleys. Although the area is just south of the Pleistocene (Illinoian) glacial limit, both the Ohio and Wabash Rivers served as major outlets for glacial meltwater and entrained sediment during glacial stages. Rapid accumulation of glacial outwash in the valleys and along the mouths of tributaries led to impoundment and extensive deposition of slackwater and lacustrine sediment in the tributary valleys. This lacustrine deposit has a complex and gradual transition with loess mantling adjacent uplands. The loess was primarily derived from the valley-bottom outwash. The uplands are underlain by faulted Pennsylvanian coal-bearing strata steeply dipping North to Northeast.

GEOTECHNICAL BEHAVIOR

The Quaternary deposits identified in the map area exhibit a wide range of grain size and geotechnical behaviors. Grain size distribution is one of the primary factors affecting the behavior of soils for geotechnical, hydrogeologic, and agricultural applications. The grain size distribution of unconsolidated sediments is dominantly controlled by the conditions under which the material was deposited. Low energy environments allow the deposition of fine-grained materials. High energy deposits limit deposition to only coarser grained materials. Eolian processes produce very well sorted (poorly graded) materials. Fluvial processes produce moderate sorting; colluvial processes produce poorly sorted deposits.

HAZARDS


Flooding is a nearly annual occurrence along the Ohio River. Floods in the late winter or early spring commonly inundate low-lying areas in the floodplain. Larger floods occur roughly every 10 to 20 years (e.g., 1913, 1945, 1964, 1997, 2007), and cover parts of the low terrace. The maximum flood of record in the valley was in 1937, flooding river towns throughout the valley. Only structures on the highest outwash terraces and the lacustrine terrace (Ql) were spared flood damage. The impact of flooding is reflected in land-use patterns through the area. Older homes and businesses have survived on the lacustrine and high outwash terraces, and on the highest parts of low terraces (Qot1, Qot1g, Qot1). Trailers and less expensively built homes are constructed on the low terraces. Only barns are found on the high parts of the floodplain (Qafp). The floodplain and low parts of the low terraces are dominantly left to woodlands or used for row-crop agriculture. Most livestock husbandry in the alluvial valleys has been abandoned and is now restricted to upland areas above the 10- to 20-year flood zone. The low-relief lacustrine terrace is locally very poorly drained.

SOILS

The silt soils that dominate the loess-mantled uplands are highly erodible. Soil piping and associated cover collapses are common hazards as ground water seeps through the silt and is commonly perched above flaggans. Great care must be taken during agricultural operations not to mobilize and lose this valuable resource.

SEISMICITY

The map area is proximal to the Wabash Valley Seismic Zone, the New Madrid Seismic Zone, and is within the Rough Creek - Shawneetown Fault Zone. Small to moderate earthquakes have been felt in the area relatively frequently. The significant thickness of unconsolidated sediment (locally as much as 150 feet in the regional map area) raise concerns about ground motion amplification of seismic waves and potential liquefaction. The variations in lithology and thickness between materials in different map units will likely cause different responses of these materials to seismic shaking.



DISCLAIMER

Although these data have been processed successfully on a computer system at the Kentucky Geological Survey (KGS), no warranty, expressed or implied, is made by KGS regarding the utility of the data on any other system, nor shall the act of distribution constitute any such warranty.

KGS does not guarantee this map or digital data to be free of errors or inaccuracies. Some cultural features originate from data sources other than KGS, and may not align with geologic features on this map. KGS disclaims any responsibility or liability for interpretations from this map or digital data or decisions based thereon.

The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Government.

ACKNOWLEDGMENTS

This map was generated using new field mapping and compilation of unpublished and previously published data and was funded in part by the U.S. Geological Survey National Cooperative Mapping Program under the STATEMAP Program authorized by the National Geologic Mapping Act of 1992. Grant numbers: 09HQ00003, and by the Kentucky Geological Survey.

Field mapping was completed by Scott Waninger from August 2009 to June 2010, with assistance from R. Counts (KGS).

Subsurface information was compiled from data on file at the Kentucky Geological Survey as well as data contributed by the Kentucky Transportation Cabinet and the U.S. Geological Survey and soil coring and auguring conducted for this project.

REFERENCES

Amos, D.H., 1970. Geologic map of the Blackford quadrangle, western Kentucky. U.S. Geological Survey Geologic Quadrangle Map GQ-0873, scale 1:24,000.

Solis, M.P., and Hettiger, A., 2000. Spatial database of the Sturgis quadrangle, western Kentucky. Kentucky Geological Survey, ser. 12. Digitally Vectorized Geologic Quadrangle Data (DVGQ)-273.12. Adapted from Kehn, T.M., 1975. Geologic map of the Sturgis Quadrangle Western Kentucky. U.S. Geological Survey Geologic Quadrangle Map GQ-1273, scale 1:24,000.

Faust, R.J., Barfield, G.R., and Willinger, G.A., 1980. A Compilation of Ground Water Quality Data for Kentucky. U.S. Geological Survey, Open File Report 80-685, p. 963.

Frye, J.C., Leonard, A.B., Wilman, H.B., and Glass, H.D., 1972. Geology and Paleontology of Late Pleistocene Lake Saline, Southeastern Kentucky, Illinois State Geological Survey, Circular 471, 44p.

Glenn, L.C., 1912. Coals of the Tradewater River Region; Kentucky Geological Survey, Bulletin #17, Serial #24, 73p.

Glenn, L.C., 1922. The Geology and Coals of Webster County, Kentucky Geological Survey, Series #6, Vol #5, 249p.

Heinrich, P.V., 1982. Geomorphology and Sedimentology of Pleistocene Lake Saline, Southern Illinois. M.S. Thesis University of Illinois: Urbana-Champaign, 145p.

Jacobs, E.H., 1981. Soil survey of Union and Webster Counties, Kentucky. U.S. Department of Agriculture, p.126.

Jilson, W.R., 1943. The Geology of Union County, Kentucky; The Standard Printing Company, 83p.

Kehn, T.M., 1974. Geologic map of the Dekoven and Saline Mines quadrangles Crittenden and Union Counties, Kentucky; U.S. Geological Survey Geologic Quadrangle Map GQ-1147, scale 1:24,000.

McFarlan, A.C., 1961. Geology of Kentucky; University of Kentucky; Kentucky Geological Survey, 331p.

Mull, D.S., Cushman, R.V., and Lambert, T.W., 1971. Public and Industrial Water Supplies of Kentucky, 1968-69. Kentucky Geological Survey Series 10, IC 20 p.107.

Nelson, W.J. and Lamm, D.K., 1986. Kentucky Geologic Map of the Shawneetown Quadrangle, Gallatin County Illinois. Illinois Geological Survey, Map GQ-1.

Palmer, J.E., 1976. Geologic map of the Grove Center quadrangle, Kentucky-Illinois, and part of the Shawneetown quadrangle, Kentucky. U.S. Geological Survey Geologic Quadrangle Map GQ-1314, scale 1:24,000.

Ray, L.L., 1965. Geomorphology and Quaternary geology of the Owensboro quadrangle, Indiana and Kentucky; U.S. Geological Survey Professional Paper 488, 71p.

Thies, C.V., 1929. The Geology of Henderson County, Kentucky, University of Cincinnati, 251p.

Universal Transverse Mercator projection, zone 16, North American Datum of 1927

Topographic base and cultural features are Kentucky Raster Graphics (KRG) from Kymartian by gov.kgmaps/KRG of Bordley

