KENTUCKY GEOLOGICAL SURVEY
James C. Cobb, State Geologist and Director
UNIVERSITY OF KENTUCKY, LEXINGTON

GEOLOGIC MAP SERIES 1
Series XII, 2001

GEOLOGIC MAP OF THE HARRODSBURG 30 x 60 MINUTE QUADRANGLE, CENTRAL KENTUCKY

Thomas N. Sparks, Garland R. Dever Jr., and Warren H. Anderson

Cartography by Terry D. Hounshell

Revised July 2012

SYSTEM	GROUP	FORMATION AND MEMBER	LITHOLOGY	THICKNESS IN METERS (FEET)
QUATERNARY		Holocene — Alluvium	Qal	0–27 (0–90)
		Holocene — Colluvium and landslide deposits	Qcol	0–9 (0–30)
		Pleistocene or Holocene — Terrace deposits	Qt	0–12 (0–40)
		Pliocene or Pleistocene — High-level fluvial deposits	QTf	0–27 (0–90)
		Pliocene or Pleistocene — Gravel and sand	QTg	0–12 (0–40)
TERT. or QUAT.		uncertain age — dolomite	Qd	
PENNSYLVANIAN	BREATHITT GROUP	Lower Pennsylvanian — Corbin Sandstone Member	Pgc	0–54 (0–180)
		Grundy Formation	Pg	0–72 (0–240)
		Alvy Creek Formation — Livingston Conglomerate Member	Pac	0–32 (0–105)
		Paragon Formation	Pap	0–33 (0–110)
MISSISSIPPIAN		Slade Formation — Poppin Rock Member	Msla	24–70 (80–230)
		Maddox Branch Member		
		Ramey Creek Member		
		Tygarts Creek Member		
		Mill Knob Member		
		Wario Run Member		
		Ste. Genevieve Member		
		St. Louis Member		
		St. Louis Limestone	Msl	20+ (65+)
		Salem and Warsaw Limestones — Undivided	Msw	16–20 (55–65)
		Multicaugh Mbr. — Rarrity Member	Mmr	
		Borden Formation — Wildie Member	Mb	85–132 (280–435)
		Halls Gap Member — Nada Member		
		Cowbell Member		
		Nancy Member		
		New Providence Shale Member		
DEVONIAN	Mid. and Upper Dev.	New Albany Shale	MDna	12–46 (40–150)
		Boyle Dolomite	Db	0–11 (0–35)
SILURIAN	Lower and Middle Silurian	Crab Orchard Formation — Brassfield Dolomite Member	Scb	0–45 (0–150)
		Brassfield Dolomite	MDsa	
ORDOVICIAN	Upper Ordovician	Dolyns Formation — Saluda Dolomite Member	Ds	0–49 (0–160)
		Preachersville Member		
		Rowland Member		
		Reba Member		
		Terrill Member		
		Ashlock Formation — Grant Lake Member	Oa	12–45 (40–150)
		Slings Creek Member		
		Gilbert Member		
		Tate Member		
		Calloway Creek Limestone	Occ	17–42 (55–140)
		Garrard Siltstone	Og	9–35 (30–115)
		Kope Formation	Ok	0–9 (0–30)
		Clays Ferry Formation	Ocf	24–73 (80–240)
		Lexington Limestone — Tanglewood Limestone Mbr. — Millersburg Strodes Creek Devils Hollow Members	Ok	0–27 (0–90)
		Tanglewood Limestone Mbr.	Ok	0–30 (0–100)
		Upper part — Sulphur Well Perryville Ls. Members	Osu	
		Tanglewood Limestone Mbr.	Ok	0–27 (0–90)
		Brannon Mbr.	Ok	0–9 (0–30)
		Tanglewood Limestone Mbr.	Ok	
		Lower part — Grier Limestone Logana Curdsville Limestone Members	Olg	36–59 (120–195)
		Tanglewood Limestone Mbr.	Ok	
		Tyrone Limestone	Oty	37 (120)
		Tyrone Limestone and Oregon Formation — Oregon Formation	Oto	
		High Bridge Group — Camp Nelson Limestone	Ocn	97+ (320+)

DESCRIPTION OF MAPPED UNITS

Qal ALLUVIUM—Silt, clay, sand, and gravel. Silt, partly clayey; clay, partly silty and sandy; sand, very fine- to coarse-grained, but mostly very fine to fine. Gravel composed of limestone, chert, dolomite, quartz, geodes, siltstone, quartzite, shale, and ironstone. Unit locally grades into unmapped colluvium.

Qcol COLLUVIUM AND LANDSLIDE DEPOSITS—Sand, gravel, and rubble. Consists of masses of shale and siltstone from Borden Formation, limestone blocks from Slade Formation, and sand, pebbles of quartz and chert, and cobbles of sandstone and conglomerate from Grundy Formation. Mapped locally in southeastern part of quadrangle.

Qt TERRACE DEPOSITS—Clay, silt, sand, and gravel. Clay, silty to sandy; silt, clayey to sandy; sand, very fine- to coarse-grained, but mostly very fine to fine. Gravel composed of quartz, chert, geodes, siltstone, ironstone, and sandstone, with rare limestone and quartzite.

QTf HIGH-LEVEL FLUVIAL DEPOSITS—Sand, gravel, silt, and clay. Sand mainly very fine- to fine-grained, partly clayey and silty; silt, clayey; clay, silty. Gravel composed of quartz, chert, geodes, quartzite, sandstone, silicified fossils, and ironstone. Deposits occur as isolated remnants of terraces along Kentucky River and its tributaries, and locally along Chaplin River.

QTg GRAVEL AND SAND—Unconsolidated conglomerate composed of quartz gravel in matrix of quartz sand and silt; scattered boulders and cobbles of conglomerate, and few pebbles of sandstone; partly cemented with limonite and silica; partly crossbedded. Unit probably consists of letdown deposits derived from Pennsylvanian conglomerate (Rice, 1984).

Od DOLOMITE—Finely to coarsely crystalline dolomite similar to dolomite in formations of Ordovician, Silurian, and Devonian age. Occurs in several small grabens in northern part of quadrangle.

Pgc CORBIN SANDSTONE MEMBER OF GRUNDY FORMATION—Sandstone, conglomerate, and siltstone. Sandstone is light-gray, medium- to very coarse-grained, common quartz pebbles. Forms steep cliffs in southeastern corner of quadrangle. Previously mapped as member of Lee Formation (Weir and others, 1971; Rice, 1972).

Pg GRUNDY FORMATION—Siltstone, shale, sandstone, and coal. Siltstone and shale, laminated, limonite-stained, and carbonaceous. Coals thin and unevenly distributed. Locally unconformable on Paragon and Slade Formations. Previously mapped as Breathitt Formation (Weir and others, 1971; Rice, 1972), or adjacent areas as Breathitt Formation, and renamed Grundy Formation of the Breathitt Group by Chesnut (1992). As mapped here, may contain lenses of Bee Rock Formation at base.

Pac ALVY CREEK FORMATION—Siltstone, shale, sandstone, and coal. Lithology similar to Grundy Formation above. Mapped unnamed coals now identified as Beaver Creek and Stearns No. 1 coal beds. Previously mapped as Lee Formation (Weir and others, 1971; Rice, 1972). Rests on Paragon Fm. with sharp, north and west, conformable on and interbrupes with Livingston Conglomerate Member, locally unconformable with Paragon and Slade Formations.

Pacl ALVY CREEK FORMATION, LIVINGSTON CONGLOMERATE MEMBER—Sandstone and conglomerate. Sandstone is quartzose and crossbedded, with common quartz pebbles. Forms steep cliffs and occupies south-trending paleovalley cut more than 18 m into Slade Formation (Rice, 1984). Smaller paleovalleys, probably tributaries to main paleovalley, are also present in the main paleovalley, but are not mappable at the scale of this map and are included in the Grundy Formation.

Pap PARAGON FORMATION—Unit consists of rocks assigned to Pennington Formation during U.S. Geological Survey–Kentucky Geological Survey cooperative mapping project, 1960–78 (Ettensohn and others, 1984). Shale, siltstone, sandstone, limestone, dolomite, and coal. Shale, partly silty and sandy, olive-gray to medium-gray, fossiliferous, locally carbonaceous. Sandstone, very fine- to medium-grained, sparsely fossiliferous, locally as much as 12 m thick at base of formation. Limestone, fine- to medium-grained, commonly fossiliferous. Dolomite, very finely crystalline, partly argillaceous, locally oolitic. Coal, several centimeters thick, locally at base of unit. Common erosional thinning and local removal of Paragon below pre-Breathitt unconformity.

Msla SLADE FORMATION—Unit includes rocks assigned to Newman Limestone and Rockhouse Member of Borden Formation during U.S. Geological Survey–Kentucky Geological Survey cooperative mapping project, 1960–78 (Ettensohn and others, 1984). Slade consists of nine members in ascending order: Poppin Rock, Maddox Branch, Ramey Creek, Tygarts Creek, Mill Knob, Wario Run, Ste. Genevieve, St. Louis, and Poppin Rock Members. Limestone, dolomite, and shale. Limestone, microgranular to very coarse-grained, bioclastic, oolitic, pelsolial; partly cherty, fossiliferous, argillaceous, brecciated, and crossbedded. Dolomite, very finely crystalline, partly argillaceous, locally sparsely cherty and geodiferous; dominant lithology in basal 2–3 m. Shale, mainly in partings and interbeds; forms distinct unit, as much as 1.5 m thick, at upper part of formation. Common erosional thinning of Slade below pre-Breathitt unconformity.

Msl ST. LOUIS LIMESTONE—Limestone, fine- to medium-grained, argillaceous, cherty, fossiliferous, geodiferous. Mapped in southwestern part of quadrangle.

Msw SALEM AND WARSAW LIMESTONES—Limestone and sandstone. Limestone, coarse- to very coarse-grained, crinoidal, partly silicified, partly cherty, partly crossbedded. Sandstone, fine-grained, calcareous, about 1.5 m thick, occurring 1.5–2.5 m above base of unit. Mapped in southwestern part of quadrangle.

Mb BORDEN FORMATION—Unit consists of seven members in ascending order, the New Providence Shale, Nancy, Cowbell, Halls Gap, Wildie, Nada, and Boyle Dolomite Members. Siltstone, sandstone, shale, and limestone. The siliciclastic-dominated Borden was reassigned to the overlying Slade Formation by Ettensohn and others (1984). Shale, siltstone, and limestone. Shale, clayey in basal part, grades upward into very silty shale and siltstone in middle and upper part; ironstone concretions; sparsely fossiliferous, with few lenses of fossiliferous limestone. Limestone, fine- to medium-grained, fossiliferous, in southeastern part of quadrangle, uppermost Borden, 10–24 m thick, consists of shale, siltstone, and minor amounts of fossiliferous limestone; in southwestern part, geodiferous, cherty, dolomitic siltstone, 12–21 m thick, with minor limestone.

MDna NEW ALBANY SHALE—Shale and dolomite. Shale, carbon-rich, silty, fissile, pyritic; few interbeds of clayey shale in middle and upper parts. In lower part of quadrangle, shale, dolomitic, silty, fissile, pyritic, interbedded with dolomitic carbonaceous, partly intraclastic; interfingered with black shale; apparently included with Boyle Dolomite on several 7.5-minute geologic quadrangle maps. Base contact conformable where unit overlies Boyle Dolomite; elsewhere base is unconformable.

Db BOYLE DOLOMITE—Dolomite, limestone, and sandstone. Dolomite, very finely to moderately crystalline, commonly cherty, partly fossiliferous, silty and sandy in lower part; vuggy in upper part; contains abundant solution breccia. Sandstone occurs locally at base, fine-grained, up to 1.8 m thick. Dolomite in basal New Albany Shale apparently included with Boyle Dolomite on several 7.5-minute geologic quadrangle maps. Unit pinches out locally. Base unconformable.

Scb CRAB ORCHARD FORMATION AND BRASSFIELD DOLOMITE—Mapping unit is composed of two formations, in ascending order: Brassfield Dolomite (0–7 m thick) and Crab Orchard Formation (0–41 m thick). Crab Orchard, in turn, consists of five members; in ascending order, the Plum Creek, Oldham, Lulbegrud Shale, Waco, and Estill Shale Members. Shale and dolomite. Shale, very finely to moderately crystalline, silty, partly fossiliferous, partly vuggy; petroliferous. Middle and upper parts of unit are mainly shale, with dolomitic interbeds; dolomite dominant in lower part. Common erosional thinning and local removal of unit below pre-Boyle unconformity.

Od DRAKES FORMATION—Unit consists of two members: in ascending order, the Rowland and Preachersville Members. Mudstone, dolomite, and limestone. Mudstone, dolomitic, calcitic; rarely fossiliferous. Dolomite, finely crystalline, argillaceous, sandy, fossiliferous, partly vuggy. Limestone occurs locally in eastern part and interbedded with mudstone. Limestone occurs locally in eastern part of quadrangle, in part fine- to coarse-grained, fossiliferous (Otter Creek coral bed); in part argillaceous. In southwestern part of quadrangle unit consists of three members: in ascending order, Rowland, Bardstown (limestone), and Preachersville Members. Common erosional erosional thinning and removal of Drakes below pre-Boyle unconformity.

Os ASHLOCK FORMATION—Unit consists of six members: in ascending order, the Tate, Gilbert, Slings Creek, Grant Lake, Terrill, and Reba Members. Limestone, mudstone, dolomite, shale, and siltstone. Limestone, incomparison to coarse-grained, partly argillaceous, partly silty, sparsely cherty; abundantly to sparsely fossiliferous. Mudstone, dolomitic, argillaceous, partly silty, sparsely fossiliferous. Dolomite, very finely crystalline, partly silty, partly interbedded with limestone. Siltstone locally interlayered and interfingers with limestone. Unit partly grades into dolomite in some areas adjacent to faults. Local erosional thinning of Ashlock below pre-Boyle unconformity.

Occ CALLOWAY CREEK LIMESTONE—Limestone, shale, and siltstone. Limestone, very fine- to very coarse-grained, fossiliferous. Shale in partings and seams between limestone beds. Siltstone interlayered with limestone in lower part. Unit partly altered to dolomite along fault in northern part of quadrangle.

Og GARRARD SILTSTONE—Siltstone, shale, and limestone. Siltstone, quartzose, micaceous, partly argillaceous, partly sandy, partly fossiliferous; shale, silty, calcareous, silty, clayey, sparsely fossiliferous, interbedded with siltstone. Limestone, fine- to coarse-grained, fossiliferous, interlayered with siltstone. Unit partly altered to dolomite along fault in northern part of quadrangle.

Ok KOPE FORMATION—Shale, siltstone, and limestone. Shale, clayey, calcareous, with interbedded siltstone and limestone.

Ocf CLAYS FERRY FORMATION—Limestone, shale, and siltstone. Limestone, micrograined to very coarse-grained, partly argillaceous, commonly fossiliferous; interbedded with about equal amounts of calcareous, clayey shale. Siltstone, calcareous; interbedded with limestone and shale in upper part. Unit partly dolomitized along fault in northern part of quadrangle. Tongues of Clays Ferry shale and limestone occur locally in upper part of Lexington Limestone.

Olu UPPER PART OF LEXINGTON LIMESTONE—Unit consists of five members: in ascending order, Millersburg, Strodes Creek, Devils Hollow, Sulphur Well, and Perryville Ls. Members. Limestone and shale. Limestone, micrograined to coarse-grained, abundantly to sparsely fossiliferous, sparsely cherty; in part with partings and interbeds of shale. In parts of quadrangle, unit interfingers with and grades laterally into Tanglewood Limestone Member of Lexington Limestone, and interfingers with Clays Ferry Formation.

Olt TANGLEWOOD LIMESTONE MEMBER OF LEXINGTON LIMESTONE—Limestone, micrograined to very coarse-grained, fossiliferous, commonly argillaceous, commonly phosphatic, sparsely to abundantly fossiliferous, rarely cherty; partly crossbedded; rarely with shale partings. In parts of quadrangle, unit interfingers with and grades laterally into upper and lower parts of the Lexington, and interfingers with Clays Ferry Formation.

Olb BRANNON MEMBER OF LEXINGTON LIMESTONE—Limestone and shale. Limestone, micrograined to fine-grained, argillaceous, silty, partly cherty, fossiliferous; interbedded with shale. Limestone, dolomitic, argillaceous, pinches out in parts of quadrangle; locally interfingers with Tanglewood Limestone Member.

Olg LOWER PART OF LEXINGTON LIMESTONE—Unit consists of three members; in ascending order, Curdsville Limestone, Logana, and Grier Limestone Members. Limestone and shale. Limestone, micrograined to very coarse-grained, fossiliferous, partly argillaceous, partly cherty; mainly irregular bedded, with shale partings and interbeds of shale. In parts of quadrangle, unit interbedded shale; in parts of quadrangle, upper part of unit interfingers with Tanglewood Limestone Member.

Oty TYRONE LIMESTONE AND OREGON FORMATION—Mapping unit is composed of two formations; in ascending order, Oregon Formation (2–20 m thick) and Tyrone Limestone (17–33 m thick). Limestone, dolomite, shale, chert, and bentonite. Tyrone Limestone consists of limestone, micrograined, partly to very coarse-grained, partly cherty, partly argillaceous, partly fossiliferous; dolomite, very finely crystalline, partly mottled with irregular bodies and seams of calcareous dolomite; few beds of pelletal and bioclastic limestone; with interbedded shale, partly dolomitic, silty, fissile, partly fossiliferous. Oregon Formation, about 7 and 24 m thick at top of Tyrone and locally at top, Oregon Formation consists of dolomite, microcrystalline to moderately crystalline, calcareous, rarely fossiliferous, partly interbedded with limestone; in parts of quadrangle, lower part of Tyrone intervenes with upper part of Oregon.

Ocn CAMP NELSON LIMESTONE—Limestone, dolomite, and shale. Limestone, micrograined, mottled with irregular bodies of very finely to moderately crystalline, calcareous dolomite; partly cherty, partly fossiliferous, commonly containing scattered biota and stringers of crystalline calcite; few interbeds of argillaceous limestone and calcareous shale, mainly in upper part. Base of formation not exposed.

af ARTIFICIAL FILL—Compacted rock debris from highway or railroad construction, and mine tailings.

EXPLANATION

- Interstate highway or parkway
- U.S. highway
- State highway
- County boundary
- City boundary
- Normal fault (U, upthrown side; D, downthrown side)
- Concealed fault
- Structure contour, feet
- Concealed contact
- Active stone quarry or mine
- Abandoned stone quarry or mine
- Abandoned sand, gravel, or clay pit
- Mineral vein
- Mineral prospect or outcrop
- Abandoned vein-mineral mine shaft

Locations of the 30 x 60 minute quadrangles covering Kentucky. The location of the Harrodsburg quadrangle is highlighted in blue.

SCALE 1:100 000

universal transverse Mercator projection, zone 16: 1927 North American datum

TOPOGRAPHIC CONTOUR INTERVAL 20 METERS

Location map for structure contours in the Harrodsburg 30 x 60 minute quadrangle. Index gives names of each mapped horizon. The horizon boundaries are shown on the geologic map as thin red dashed lines. Contour interval is 100 feet. Datum is mean sea level. Where dip is relatively steep, contour interval is 100 feet.

Index map showing the locations of the 32 quadrangles used in the digital compilation of the Harrodsburg 30 x 60 minute quadrangle. Index also shows names of the individual 7.5-minute quadrangle maps and their USGS geologic quadrangle map (GQ) numbers.

ACKNOWLEDGMENTS: This map was digitally generated from 1:24,000-scale geologic maps under the STATEMAP Program authorized by the National Geologic Mapping Act of 1992, Grant No. 99HQAG2035, and matching funds from the Kentucky Geological Survey, National Cooperative Geologic Mapping Program.

DISCLAIMER: Although these data have been processed successfully on a computer system at the Kentucky Geological Survey, no warranty, expressed or implied, is made by KGS regarding the utility of the data on any other system, nor shall the act of distribution constitute any such warranty. The KGS does not warrant this map or digital data to be free of errors or inaccuracies. KGS disclaims any responsibility or liability for interpretations from this map or digital data, or decisions based thereon.

The views and conclusions contained in this document are those of the authors and should not be interpreted as representing opinions or policies of the sponsor, either expressed or implied, of the U.S. Government.

For information on obtaining copies of this map and other Kentucky Geological Survey maps and publications call:

Publication Sales
(859) 257-3896

View the KGS World Wide Web site at
www.uky.edu/KGS/

NOTE: cross section is diagrammatic

NW — A — A' — SE

vertical exaggeration 6x

CINCINNATI ARCH

JESSAMINE DOME

LEXINGTON FAULT SYSTEM

IRVINE-PAINT CREEK FAULT SYSTEM

Kentucky River

Jessamine Creek

Hickman Creek

400 m 1312 ft
200 m 656 ft
(sea level) 0
–200 m –656 ft
–400 m –1312 ft

Clays Ferry Formation
Lexington Limestone
High Bridge Group
Beekmantown Dolomite (Knox Group)
Copper Ridge Dolomite (Knox Group)
Garrard Siltstone
Calloway Creek Limestone
Ashlock Formation
Drakes Formation
Crab Orchard Formation and Brassfield Dolomite
New Albany Shale
Pottsville/Cumberland Escarpment
Livingston Conglomerate
Grundy Formation
Paragon Formation
Corbin Sandstone
Slade Formation
Borden Formation

INTRODUCTION

This version of the geology of the Harrodsburg 30 x 60 minute quadrangle was digitally compiled mostly from U.S. Geological Survey 7.5-minute geologic quadrangle maps (GQ's), as cited in the references. The GQ's, in turn, were the product of mapping carried out between the U.S. Geological Survey and the Kentucky Geological Survey from 1960 to 1978. Several regional studies on mapping and stratigraphy (e.g. Cressman, 1973; Ettensohn and others, 1984; Chesnut, 1992) have been published since completion of mapping and fact correlation. These changes are shown on this map, and were necessary for compilation of regional maps and for stratigraphic continuity between quadrangles. The 7.5-minute quadrangles that make up the Harrodsburg 30 x 60 minute quadrangle are shown in the index map.

PROCEDURE

The 7.5-minute quadrangle maps were digitally compiled using a semi-automated data-capture technique to convert hard-copy geologic maps into digital format. Compiling 7.5-minute maps into a 30 x 60 minute map required the resolution of significant problems, such as (1) correlating geologic contacts across quadrangle boundaries, (2) resolving nonuniform structure-contour intervals, (3) digitally joining geologic contacts and Quaternary alluvium boundaries and inferred contacts. The metadata portion of the digital file provides detailed information about the conversion process. Formation codes were assigned using the American Association of Petroleum Geologists' Publication Format Committee (Cohee, 1967), which was modified by the Kentucky Geological Survey for state-specific use. Formations and contact-line boundaries were not mapped the same way on each of the 7.5-minute quadrangle maps, since they were compiled between 1960 and 1978. Resolution of the differences between quadrangles was necessary for digital compilation into a geographic information system (GIS) environment. In addition, numerous small members mapped on individual 7.5-minute quadrangle maps are too small to be mapped at a scale of 1:100,000 on a 30 x 60 minute quadrangle map. These problems were resolved by applying geologic, cartographic, and GIS standards appropriate for the scale of the map.

This map is the result of more than 30 years of geological-geologic field work took place. When there were problems in stratigraphic correlation between quadrangles, the best current data available were used to resolve these differences.

GEOLOGIC SETTING AND STRUCTURAL GEOLOGY

The geology of the Harrodsburg 30 x 60 minute quadrangle mainly consists of flat-lying sedimentary rocks of Mississippian and Pennsylvanian age. These rocks extend across the north-trending Cincinnati Arch, a structural feature that extends from Ohio through central Kentucky to Nashville. The Jessamine Dome, a regional domal structure, lies along the Cincinnati Arch in the southeastern part of the quadrangle. Most of the Jessamine Dome occurs in the Harrodsburg quadrangle in southern Jessamine County.

The apex is reflected in the circular outcrop pattern of the Upper Ordovician units in the center of the map. Three major fault systems form the boundaries of the Jessamine Dome: (1) the Lexington Fault System, which bisects the quadrangle from north to south, (2) the Kentucky River Fault System, which trends northeast-southwest in the northwest part of the quadrangle and represents the northern boundary of the Rome Trough, and (3) the Irvine–Paint Creek Fault System, which is oriented east-northeast across the southeastern part of the quadrangle. The western boundary of the Irvine–Paint Creek Fault System is the eastern limb of the Jessamine Dome. Detailed investigations of the Jessamine Dome, Rome Trough, and Irvine–Paint Creek Fault System are summarized by Cressman (1963), Harris and Drahovzal (1996), and Drahovzal and Noger (1995).

ECONOMIC AND ENGINEERING GEOLOGY

Limestone, vein minerals, gas, clay, and coal are the principal mineral resources in the Harrodsburg 30 x 60 minute quadrangle. Limestone suitable for construction aggregate and agricultural products is being quarried from the Lexington Limestone in Boyle and Mercer Counties and mined from the High Bridge Group in Fayette, Jessamine, Garrard, and Madison Counties (Anderson and others, 1982). Limestone was formerly mined in the Harrodsburg quadrangle, and many have been mined for barite, fluorite, calcite, and sphalerite (Anderson and others, 1982). Fluorite was mined at several sites near Mudcreek Landing in southern Woodford County during World War I. Barite was mined in Jessamine County during World War II. The Chinn calcite vein and other veins were mined for fluorite in Mercer County during this same period. A vein was mined for barite in Lincoln County in the 1960's, and another vein was mined for sphalerite at the Lexington Quarry Company in Jessamine County during the 1980's. These veins are currently considered to be prospects. Core drilling for several thousand feet along strike. Several stratigraphic tests have been drilled in search of similar mineral occurrences and petroleum in the subsurface of the quadrangle. Traces of galena, sphalerite, fluorite, and barite mineralization have been noted at depth in the Knox Group, but these deposits are not economically developed.

Several oil and gas fields have been discovered along the previously mentioned fault systems (Wilson and Sutton, 1976) and other anticlinal structures may play an important role in the accumulation of petroleum resources, particularly Cambrian natural gas. The following fields have produced oil from shallow reservoirs: Bryantsville, Boyle County; Lancaster, Garrard County; Carlisville, Garrard County; Lanhart, Jackson County; Junks, Estill County; Neals Creek and Jumbo Fields, Lincoln County. Cumulative production from the St. Peter Sandstone in the Trapp Gas Field, Clark County, is about 400 million cubic feet (MMCF). Several deep gas wells have been drilled in the Cambrian, and had a show of gas, but none have been developed to supply a pipeline to market the gas. For more information on the geology and production in these fields, see Rosen and Walker (1999).

Some unnamed coal beds occur in the southeastern part of the quadrangle, but they are discontinuous and have not been extensively mined. Sand, gravel, and clay are mined from the High Bridge Group and from additional deposits in the eastern and southwestern parts of the quadrangle, but most operations are currently inactive.

Clay from the Borden Formation could have commercial uses, although it is not currently mined. Analysis of samples from the southern part of the quadrangle indicates potential for industrial uses in the pottery, brick, and tile industry and for use as lightweight aggregate. Shale from the Borden and Slade Formations alone may also have use as flagstone and aggregate.

Slope instability in thick sequences of shales of the Borden Formation affects construction and engineering projects. Landslides or slumps could occur when developed slopes in the Borden are saturated with water.

HYDROGEOLOGY

The Harrodsburg quadrangle covers all of regions 2, 3, and 4 (Palmquist and Hall, 1980a, b, c) of the Kentucky River Basin watershed (Carey and others, 1964). The availability and quality of ground water in this area is discussed in Hall and Palmquist (1960), Palmquist and Hall (1960,b), Carey (1992), Kipp (1997), and Currens and Ray (1998). Most ground water is obtained from wells less than 100 feet deep in shallow alluvial or bedrock aquifers. Wells in the carbonate rock aquifers may provide adequate water for farms or commercial operations. Shallow carbonate aquifers in the quadrangle are karstic. Ground water in karst aquifers occurs in solution-enlarged joints and bedding planes and in caves. Sinkholes are the most favorable locations for water-well drilling, so several dry holes may be drilled before a well producing an adequate supply for domestic use is found. Where favorably located, springs provide a more predictable supply. Water from all wells and springs in karst aquifers should be sanitized before being used for human consumption.

Anderson, W.H., and Dever, G.R., Jr. 1996, Mineral and fuel resources map of Kentucky: Kentucky Geological Survey, ser. 11, Map and Chart Series 21, scale 1:500,000.

Anderson, W.H., Dever, G.R., Jr., and Grean, R., Sand, barite deposits of Kentucky: Kentucky Geological Survey, ser. 11, Bulletin 1, 56 p.

Carey, D.I., 1992, Water in the Kentucky River Basin: Kentucky Geological Survey, ser. 11, Information Circular 37, 96 p.

Carey, D.I., 2000, Spatial database of the Burkeye quadrangle, central Kentucky: Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-843. Adapted from Wolcott, D.E., 1970, Geologic map of the Burkeye quadrangle, central Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-844, scale 1:24,000.

Carey, D.I., Hettinger, C.P., Dever, G. Kipp, J.A., Wunsch, R.S., and Conrad, D., 1994, Ground-water in the Kentucky River Basin: Kentucky Geological Survey, ser. 11, Information Circular 52, 67 p.

Carey, D.I., and Hettinger, C.P., 2000, Spatial database of the Harrodsburg quadrangle, central Kentucky: Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-945. Adapted from Wolcott, D.E., and Cressman, E.R., 1971, Geologic map of the Harrodsburg quadrangle, central Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-946, scale 1:24,000.

Chesnut, D.R., Jr., 1992, Stratigraphic and structural framework of the Carboniferous rocks of the central Appalachian Basin in Kentucky: Kentucky Geological Survey, ser. 11, Bulletin 3, 42 p.

Cissak, E.A., 2000a, Spatial database of the Little Hickman quadrangle, Jessamine and Fayette Counties, Kentucky: Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-792. Adapted from Wolcott, D.E., 1969, Geologic map of the Little Hickman quadrangle, central Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-800, scale 1:24,000.

Cissak, E.A., 2000b, Spatial database of the Lancaster quadrangle, Lincoln and Garrard Counties, Kentucky: Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-846. Adapted from Weir, G.W., 1971, Geologic map of the Lancaster quadrangle, central Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-885, scale 1:24,000.

Cissak, E.A., 2000c, Spatial database of the Nicholasville quadrangle, Jessamine and Fayette Counties, Kentucky: Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-767. Adapted from MacQueen, W.C., Jr., 1968, Geologic map of the Nicholasville quadrangle, central Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-767, scale 1:24,000.

Cissak, E.A., 2000d, Spatial database of the Paint Lick quadrangle, east-central Kentucky: Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-840. Adapted from Weir, G.W., 1965, Geologic map of the Paint Lick quadrangle, east-central Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-470, scale 1:24,000.

Cissak, E.A., 2000e, Spatial database of the Wilmore quadrangle, Madison and Fayette Counties, Kentucky: Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-847. Adapted from Simmons, E.R., and Hrabar, S.V., Jr., 1967, Geologic map of the Wilmore quadrangle, central Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-647, scale 1:24,000.

Cohee, G.V., chairman, 1967, Standard stratigraphic code adopted by the American Association of Petroleum Geologists: American Association of Petroleum Geologists Bulletin v. 51, p. 2146–2151.

Cressman, E.R., 1973, Lithostratigraphy and depositional environments of the Lexington Limestone (Middle Ordovician) in central Kentucky: U.S. Geological Survey Professional Paper 768, 61 p.

Currens, J.C., and Ray, J.A., 1998, Mapped karst ground water basins in the Harrodsburg 30 x 60 minute quadrangle: Kentucky Geological Survey, ser. 11, Map and Chart Series 18, scale 1:100,000.

Drahovzal, J.A., and Noger, M.C., 1995, Preliminary map of the structure of the Precambrian surface in eastern Kentucky: Kentucky Geological Survey, ser. 11, Map and Chart Series 15, scale 1:500,000.

Ettensohn, F.R., Rice, C.L., Dever, G.R., Jr., and Chesnut, D.R., 1984, Slade and Paragon Formations–New stratigraphic nomenclature for Mississippian rocks along the Cumberland Escarpment in Kentucky: U.S. Geological Survey Bulletin 1605-B, 37 p.

Greene, R.C. 1960, Geologic map of the Perryville quadrangle, Estill and Madison Counties, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-998, scale 1:24,000.

Hall, F.R., and Palmquist, W.N., Jr., 1960, Availability of ground water in Clark, Estill, Madison, and Powell Counties, Kentucky: U.S. Geological Survey Hydrologic Investigations Atlas HA-25, scale 1:125,000.

Harris, D.C., and Drahovzal, J.A., 1996, Cambrian hydrocarbon potential indicated in Kentucky's Rome Trough: Kentucky Geological Survey, ser. 11, Information Circular 54, 11 p.

Hettinger, C.P., 2000, Spatial database of the Stanford quadrangle, Lincoln and Boyle Counties, Kentucky: Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-1137. Adapted from Shawe, F.R., and Wigley, P.B., 1974, Geologic map of the Stanford quadrangle, Boyle and Lincoln Counties, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1137, scale 1:24,000.

Kipp, J.A., 1997, Fresh-water aquifer in the Knox Group (Cambrian–Ordovician) of central Kentucky: Kentucky Geological Survey, ser. 11, Report of Investigations 12, 36 p.

McGrain, P., Kessler, T.A., and Tesker, T.C., 1960, Miscellaneous clay and shale analyses for 1957–1959: Kentucky Geological Survey, ser. 10, Report of Investigations 3, 51 p.

Mills, J.M., 2000, Spatial database of the Moberly quadrangle, Jessamine and Fayette Counties, Kentucky: Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-664. Adapted from Greene, R.C., 1968, Geologic map of the Moberly quadrangle, Jessamine and Fayette Counties, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-664, scale 1:24,000.

Miller, J.M., 2000a, Spatial database of the Parksville quadrangle, Estill and Madison Counties, Kentucky: Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-690. Adapted from Greene, R.C., 1960, Geologic map of the Parksville quadrangle, Estill and Madison Counties, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-998, scale 1:24,000.

Miller, J.M., 2000b, Spatial database of the Perryville quadrangle, Mercer and Boyle Counties, Kentucky: Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-1165. Adapted from Cressman, E.R., 1975, Geologic map of the Perryville quadrangle, Mercer and Boyle Counties, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1165, scale 1:24,000.

Sullivan, J.M., 2000, Spatial database of the Palmer quadrangle, east-central Kentucky: Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-813. Adapted from Simmons, Q.C., 1967, Geologic map of the Palmer quadrangle, east-central Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-813, scale 1:24,000.

Thompson, M.F., 2000b, Spatial database of the Cornishville quadrangle, Mercer County, Kentucky: Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-1075. Adapted from Cressman, E.R., 1975, Geologic map of the Cornishville quadrangle, Mercer County, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1075, scale 1:24,000.

Thompson, M.F., 2000c, Spatial database of the Keene quadrangle, Jessamine and Woodford Counties, Kentucky: Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-440. Adapted from Cressman, E.R., 1965, Geologic map of the Keene quadrangle, central Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-440, scale