

**MAPPED KARST GROUND-WATER BASINS IN THE
BEAVER DAM 30 x 60 MINUTE QUADRANGLE**

Joseph A. Ray
Kentucky Natural Resources and Environmental Protection Cabinet—
Division of Water
James C. Currens
Kentucky Geological Survey

LEGEND

- Area of potential karst ground-water basin development
- Area of limited karst ground-water basin development
- Inferred perennial ground-water flow route
- Subsurface overflow (high-flow) route
- Surface overflow (high-flow) route
- Ground-water basin catchment boundary
- Intermittent lake
- Stream sink or swallet
- Underflow spring (perennial)
- Overflow spring (high flow)
- Karst window or sinking spring
- Cave stream
- Other tracer-injection point
- Water well
- Kentucky Division of Water AKGWA spring identification number

EXPLANATION


This map shows karst ground-water basins in the Beaver Dam quadrangle, determined primarily by ground-water tracer studies. It can be used to quickly identify the ground-water basins and springs to which a site may drain. Major springs and the relative size of their catchment areas can be evaluated for potential as water supplies. The map also serves as a geographic index to literature on karst ground water in the area.

This map is designed for regional and preliminary hydrologic investigations. Features such as springs and swallets are much too small to precisely locate on this map with a scale small enough to show regional relationships. The user is referred to the literature for detailed site descriptions. The data used to compile this map were obtained by numerous investigators over the last 25 years. The underflow spring draining a ground-water basin is assigned a unique identification number, referred to as the AKGWA number (Assembled Kentucky Ground Water Database). Individual basins are identified by the underflow spring name and AKGWA number. The authors of tracer data are identified by number in the "Data Source" column of the key, and are listed in "References Cited" in order of publication or research date.

Although ground-water flow routes shown here have been established by tracer studies, with the exception of mapped cave streams, the precise flow paths are unknown and are inferred or interpreted using water-level data, geologic structure, or surface features. Arrows show the direction of ground-water flow and tracer recovery locations. Conduit flow is illustrated as either thick trunk-flow lines or thin tributary-flow lines. The locations of some ground-water basins are inferred, based on the existence of a significant spring system and the delineation of adjacent basins. The position of ground-water basin boundaries should be considered approximate because of the map's scale and because boundaries can shift during high-water conditions. Also, excess flow may exit or enter a basin via surface or subsurface overflow routes. Additional overflow routes probably exist. Although most of the results of ground-water tracing shown on this map were obtained during moderate- or high-flow conditions, the ground-water basins are illustrated in base flow because base flow is the most common flow condition. The main spring draining the basin is assumed to be an underflow spring that preferentially drains base flow. Overflow springs discharge during high flow. Generally, names of ground-water basins are derived from these main springs. Not all additional springs are shown because of the small map scale.

DISCLAIMER: This map is subject to revision upon receipt of new hydrologic data. The unshaded area (shown in white on the map) is karst. The shaded area (shown in light brown) is largely underlain by noncarbonate rocks and has minimal development of karst. Karst features are only shown in those areas where tracer tests have been conducted. The user should consult the "References Cited" for additional information.

¹Worthington, S.R.H., 1991, Karst hydrogeology of the Canadian Rocky Mountains: Hamilton, Ontario, McMaster University, Ph.D. dissertation, 380 p.



KEY

| AKGWA No. | Spring Name | Data Source |
|-----------|------------------------|---------------------------------|
| 0050 | Double | (12) |
| 0051 | Graham | (2, 3, 5, 6, 7, 8, 9, 15, 26) |
| 0062 | Pike | (1, 6, 7, 9, 15) |
| 0083 | Turnhole | (2, 3, 6, 7, 9, 15, 17, 18, 24) |
| 0186 | Big | (17, 25) |
| 0191 | Echo River | (1, 6, 7, 9, 15, 17) |
| 0193 | River Stix | (1, 6, 7, 9, 15, 17) |
| 0195 | Buffalo Creek | (4, 16, 17, 19, 20, 22) |
| 0197 | Mile 205.7 | (14, 16, 17, 22) |
| 0198 | Gristle Mill | (9, 15) |
| 0199 | Sandhouse Cave | (6, 9, 15) |
| 0200 | Sand Cave | (9, 15) |
| 0202 | Maunum | (13) |
| 0228 | Wellb Cave/Bacon Creek | (9, 15) |
| 0230 | Bell | (10) |
| 0231 | Goodman | (10) |
| 0232 | Suda | (9, 15) |
| 0235 | Buck Island | (9, 15) |
| 0236 | Copeland | (10) |
| 0238 | Twilight | (10) |
| 0261 | Britton | (10) |
| 0262 | Avery | (10) |
| 0263 | Greencastle Big | (9, 15) |
| 0264 | Sharp Bluehole | (9, 15) |
| 0265 | Cab Run Complex | (17) |
| 0266 | Mile 5 | (17) |
| 0271 | Running Branch | (17) |
| 0272 | Wilson Cave Hole | (17) |
| 0276 | Doye Ford | (17) |
| 0278 | Ligt Creek | (17) |
| 0279 | Nappers Notch | (14, 17) |
| 0280 | Marin | (9, 15, 26) |
| 0533 | Martin | (9, 15, 26) |
| 0535 | Posthouse | (9, 15, 26) |
| 0536 | McCoy Bluehole | (6, 9, 10, 15, 17) |
| 0702 | Westbrook Creek | (21) |
| 1248 | Lock Bend | (17) |
| 1249 | Sullivan Hollow | (17) |
| 1431 | Sail Hollow | (17) |
| 1432 | Hillbender | (17) |
| 1433 | First Creek | (17) |
| 1434 | Watworks | (17) |
| 1435 | Cotton Gin Hollow | (9, 15) |
| 1436 | Pratt Sulphate Cave | (9, 15) |
| 1437 | Good Lake | (9, 15) |
| 1438 | Indiah Creek | (17) |
| 1439 | Whittle | (17) |
| 1440 | Dry Branch | (27) |
| 1441 | Tribstone | (17) |
| 1442 | Pansy Hollow | (17) |
| 1443 | Creefish | (17) |
| 1444 | Sycamore | (17) |

REFERENCES CITED

(1) Anderson, R.B., 1925, Investigation of a proposed dam site in the vicinity of Mammoth Cave, Kentucky, cited in Brown, R.F., 1966, Hydrology of the cavernous limestone of the Mammoth Cave area, Kentucky: U.S. Geological Survey Water-Supply Paper 1837, 64 p.

(2) Miotke, F.D., and Pappeberg, Hans, 1972, Geomorphology and hydrology of the Sinkhole Plain and Glasgow Upland, central Kentucky karst: Caves and Karst, v. 14, no. 4, p. 25-32.

(3) Wells, S.G., 1973, Geomorphology of the Sinkhole Plain in the Pennsylvanian Plateau of the central Kentucky karst: Cincinnati, Ohio, University of Cincinnati, Master's thesis, 108 p.

(4) Quinlan, J.F., 1974, Unpublished ground-water tracing data: Mammoth Cave National Park, Kentucky.

(5) Magerlein, S.D., and Dillon, Clarence, 1976, Unpublished spring diving data (12-28-76): Graham Springs Complex, Kentucky.

(6) Quinlan, J.F., and Rowe, D.R., 1977, Hydrology and water quality in the central Kentucky karst: Phase I: University of Kentucky, Kentucky Water Resources Research Institute, Research Report 101, 93 p.

(7) Quinlan, J.F., and Rowe, D.R., 1978, Hydrology and water quality in the central Kentucky karst: Phase II, part A: Preliminary summary of the hydrogeology of the Mill Hole sub-basin of the Turnhole Spring ground-water basin: University of Kentucky, Kentucky Water Resources Research Institute, Research Report 109, 42 p.

(8) Powell, R.L., 1979, Unpublished ground-water tracing data: Warren County, Kentucky.

(9) Quinlan, J.F., and Ray, J.A., 1981, Groundwater basins in the Mammoth Cave region, Kentucky, showing springs, major caves, flow routes, and potentiometric surface: Friends of the Karst, Occasional Publication 1, scale 1:38,000.

(10) Quinlan, J.F., and Ray, J.A., 1981-85, Unpublished ground-water tracing data: Mammoth Cave National Park, Kentucky.

(11) Saunders, J.W., 1984, Pratt Sulphate Cave, Warren County, Kentucky: Central Kentucky Cave Survey, Bulletin 1, p. 208-209.

(12) Schindler, G.M., 1984, Enteric contamination of an urban karstified carbonate aquifer: The Double Springs drainage basin, Bowling Green, Kentucky: Bowling Green, Western Kentucky University, Master's thesis, 147 p.

(13) Quinlan, J.F., 1986, Application of dye-tracing to date-site evaluation in a Kentucky karst area, U.S.A., International Symposium on Karst Water Resources, Ankara, Turkey, Proceedings, unpaginated.

(14) Quinlan, J.F., and Schaftall, Tim, 1988, Unpublished ground-water tracing data: Mammoth Cave National Park, Kentucky.

(15) Quinlan, J.F., and Ray, J.A., 1988, Groundwater basins in the Mammoth Cave region, Kentucky, showing springs, major caves, flow routes, and potentiometric surface: Friends of the Karst, Occasional Publication 2, scale 1:38,000.

(16) Quinlan, J.F., and Ray, J.A., 1988, Unpublished ground-water tracing data: Mammoth Cave National Park, Kentucky.

(17) Meiman, Joe, and Ryan, M.T., 1990-94, Unpublished ground-water tracing data: Mammoth Cave National Park, Kentucky.

(18) Murphy, J.D., 1992, Determining the hydrology of the Turnhole Spring ground-water basin of Mammoth Cave National Park, Kentucky, using quantitative dye tracing: Durham, England, University of Durham, undergraduate dissertation, 51 p.

(19) Hammon, D.L., 1992, Hydrology and geology of the Buffalo Creek area on the north side of Mammoth Cave National Park, Kentucky: Eastern Kentucky University, Master's thesis, 70 p.

(20) Ryan, M.T., 1987, Using newly-developed quantitative dye tracing techniques to determine the karst hydrology of the Buffalo Spring ground-water basin of Mammoth Cave National Park, Kentucky: Richmond, Eastern Kentucky University, Master's thesis, 121 p.

(21) Smith, J.H., and Crawford, N.C., 1989, Groundwater flow in the vicinity of the Curtis Ray Landfill, Warren County, Kentucky: Bowling Green, Ky., Unpublished consultant's report.

(22) Uhlenschuck, C.R., 1993, Hydrogeology and groundwater quality of the Maple Springs area, Mammoth Cave National Park, Kentucky: Richmond, Eastern Kentucky University, master's thesis, 138 p.

(23) Crawford, N.C., 1994, Unpublished ground-water tracing data: Bowling Green, Ky., Crawford and Associates.

(24) Meiman, Joe, and Capps, A.S., 1995, Unpublished ground-water tracing data: Mammoth Cave National Park, Kentucky.

(25) Ryan, M.T., and Meiman, Joe, 1996, An examination of short-term variations in water quality at a karst spring in Kentucky: Ground Water, v. 34, no. 1, p. 23-30.

(26) Ray, J.A., Ewers, R.O., and Ibsen, P.J., 1997, Mapping water-supply protection areas for leaky, perched karst groundwater systems, in Beck, B.F., ed., 6th Multidisciplinary Conference on Sinkholes and Engineering and Environmental Impacts of Karst, Springfield, Mo., Proceedings, p. 153-156.


(27) Meiman, Joe, 1997, Unpublished ground-water tracing data: Mammoth Cave National Park, Kentucky.

September 1998

CAUTION: Prolonged exposure to sunlight or contact with water will damage this photo.

Base map compiled from U.S. Geological Survey digital line graphs.

ACKNOWLEDGMENTS:
We thank the many karst investigators who have contributed data for this map. Without their cooperation this map would not have been possible.



Location of the 1:100,000-scale quadrangle shown on Kentucky. This map is the Beaver Dam quadrangle, in light brown in green.

For information on obtaining copies of this map and other Kentucky Geological Survey maps and publications call:
Publications Sales office
(606) 257-3905
Visit the KGS World Wide Web site at <http://www.uky.edu/KGS/home.htm>

Cartography by Terry Houshall