

**MAPPED KARST GROUND-WATER BASINS IN THE BOWLING GREEN 30x60 MINUTE QUADRANGLE**

Joseph A. Ray  
Kentucky Natural Resources and Environmental Protection Cabinet—  
Division of Water  
James C. Currens  
Kentucky Geological Survey

**EXPLANATION**

This map shows karst ground-water basins in the Bowling Green quadrangle, determined primarily by ground-water tracer studies. It can be used to quickly identify the ground-water basins and springs to which a site may drain. Major springs and the relative size of their catchment areas can be evaluated for potential as water supplies. The map also serves as a geographic index to literature on karst ground water in the area.

This map is designed for regional and preliminary hydrologic investigations. Features such as springs and swallets are much too small to precisely locate on this map with a scale small enough to show regional relationships. The user is referred to the literature for detailed site descriptions. The data used to compile this map were obtained by numerous investigators over the last 25 years. The underflow spring draining a ground-water basin is assigned a unique identification number, referred to as the AKGWA number (Assembled Kentucky Ground Water Database). Individual basins are identified by the underflow spring name and AKGWA number. The authors of tracer data are identified by number in the "Data Source" column of the key, and are listed in "References Cited" in order of publication or research date.


Although ground-water flow routes shown here have been established by tracer studies, with the exception of mapped cave streams, the precise flow paths are unknown and are inferred or interpreted using water-level data, geologic structure, or surface features. Arrows show the direction of ground-water flow and tracer recovery locations. Conduit flow is illustrated as either thick trunk-flow lines or thin tributary-flow lines. The locations of some ground-water basins are inferred, based on the existence of a significant spring system and the delineation of adjacent basins. The position of ground-water basin boundaries should be considered approximate because of the map's scale and because boundaries can shift during high-water conditions. Also, excess flow may exit or enter a basin via surface or subsurface overflow routes. Additional overflow routes probably exist. Although most of the results of ground-water tracing shown on this map were obtained during moderate- or high-flow conditions, the ground-water basins are illustrated in base flow because base flow is the most common flow condition. The main spring draining the basin is assumed to be an underflow spring that preferentially drains base flow. Overflow springs discharge during high flow. Generally, names of ground-water basins are derived from these main springs. Not all additional springs are shown because of the small map scale.

**DISCLAIMER:** This map is subject to revision upon receipt of new hydrologic data. The unshaded area (shown in white on the map) is karst. The shaded area (shown in light brown) is largely underlain by noncarbonate rocks and has minimal development of karst. Karst features are only shown in those areas where tracer tests have been conducted. The user should consult the "References Cited" for additional information.

Worthington, S.R.H., 1991. Karst hydrogeology of the Canadian Rocky Mountains: Hamilton, Ontario, McMaster University, Ph.D. dissertation, 380 p.

**LEGEND**

- Area of potential karst ground-water basin development
- Area of limited karst ground-water basin development
- Inferred perennial ground-water flow route
- Subsurface overflow (high-flow) route
- Surface overflow (high-flow) route
- - - Ground-water basin catchment boundary
- - - Ground-water basin catchment sub-boundary
- Intermittent lake
- Stream sink or swallet
- Underflow spring (perennial)
- Overflow spring (high flow)
- Karst window or sinking spring
- Cave stream
- Other tracer-injection point
- Water well
- Kentucky Division of Water AKGWA spring identification number
- Spring name




Base map compiled from U.S. Geological Survey digital line graphs. CAUTION: Prolonged exposure to sunlight or contact with water will damage this map.

- (1) Mieske, F.D., and Popenberg, H., 1972. Geomorphology and hydrology of the Sinkhole Plain and Glasgow Upland, central Kentucky karst: Caves and Karst, v. 14, no. 4, p. 25-32.
- (2) George, A.I., 1973. Unpublished ground-water tracing data: Louisville, Ky., A.I. George Consultants.
- (3) Wells, S.G., 1973. Geomorphology of the Sinkhole Plain in the Pennyrival Plateau of the central Kentucky karst: Cincinnati, Ohio, University of Cincinnati, master's thesis, 108 p.
- (4) Connor, D.G., 1976. The lower reaches of Long Creek, Kentucky: Karst anomaly in Allen County, Bowling Green, Western Kentucky University, master's thesis, 146 p.
- (5) Quinlan, J.F., and Rowe, D.R., 1977. Hydrology and water quality in the central Kentucky karst: Phase I: University of Kentucky, Water Resources Research Institute, Research Report 101, 93 p.
- (6) Quinlan, J.F., and Rowe, D.R., 1978. Hydrology and water quality in the central Kentucky karst: Phase II, part A: Preliminary summary of the hydrogeology of the Mill Hole sub-basin of the Turnhole Spring groundwater basin: University of Kentucky, Water Resources Research Institute, Research Report 109, 42 p.
- (7) Mlynek, J.E., 1979. Annual report: Western Kentucky speleological survey: Murray State University, 84 p.
- (8) Cabbage, J.C., 1981. The hydrology of the Sinking Creek system, Logan and Simpson Counties, Kentucky: Bowling Green, Western Kentucky University, master's thesis, 83 p.
- (9) Quinlan, J.F., and Ray, J.A., 1981. Groundwater basins in the Mammoth Cave region, Kentucky, showing springs, major caves, flow routes, and potentiometric surface: Friends of the Karst, Occasional Publication 1, scale 1:138,000.
- (10) Quinlan, J.F., 1982. Interpretation of dye-tests and related studies, Caldwell Lace Leather Co., waste disposal sites, Logan Co., Kentucky: Mammoth Cave, Ky., unpublished consultant's report, 35 p.
- (11) Ewers, R.O., 1983. Groundwater flow directions in the vicinity of Caldwell Lace Leather Co., waste disposal sites, Richmond, Ky., unpublished consultant's report, 9 p.
- (12) Schmidt, G.M., 1984. Enteric contamination of an urban karstified carbonate aquifer: The Double Springs drainage basin, Bowling Green, Kentucky: Bowling Green, Western Kentucky University, master's thesis, 147 p.
- (13) Able, A.S., 1984. Subsurface drainage systems west of Bowling Green, Warren County, Kentucky: Unpublished ground-water tracing report: Bowling Green, Western Kentucky University.
- (14) Able, A.S., 1985. Subsurface drainage systems of Brush Creek and Clear Fork Creek of Warren County, Kentucky: Unpublished ground-water tracing report: Bowling Green, Western Kentucky University.
- (15) Arruda, J.K., 1985. Dye traces in the Mt. Ayr, Briarwood Manor Estates area: Unpublished ground-water tracing report: Bowling Green, Western Kentucky University.
- (16) Crawford, N.C., 1985a. Groundwater flow and geologic structure, Lost River groundwater basin, Western Kentucky University, Center for Cave and Karst Studies, scale 1:57,600.
- (17) Crawford, N.C., 1985b. Groundwater flow in the vicinity of Kentucky Stone Company, Simpson County, Kentucky: Bowling Green, Ky., Crawford and Associates, 6 p.
- (18) Franzenfelder, A., and Smith, D., 1985. Karst hydrologic study of the Woodburn, Kentucky, area: Unpublished ground-water tracing report: Bowling Green, Western Kentucky University.
- (19) Groves, C.G., 1985. Karst hydrogeology of the Sunken Spring drainage basin, Warren County, Kentucky: Unpublished ground-water tracing report: Bowling Green, Western Kentucky University.
- (20) Hill, W.M., and Lopez, L.G., 1985. Subsurface drainage, Rockfield area, Kentucky: Unpublished ground-water tracing report: Bowling Green, Western Kentucky University.
- (21) Quarles, M., and Beaven, A., 1985. Karst hydrologic features of the Plano, Kentucky, area: Unpublished ground-water tracing report: Bowling Green, Western Kentucky University.
- (22) Arruda, J.K., 1986. Groundwater flow routes of north Bowling Green, Kentucky: Unpublished ground-water tracing report: Bowling Green, Western Kentucky University.
- (23) Butorak, K., and Lassaline, E., 1986. Subsurface drainage, Petrow Browning area, Kentucky: Unpublished ground-water tracing report: Bowling Green, Western Kentucky University.
- (24) Dyas, M., and Foreyhe, P., 1988. Wilson-Stanley cave system: Central Kentucky Cave Survey, Bulletin 2, p. 143-144.
- (25) Crawford, N.C., 1989. Unpublished ground-water tracing data: Bowling Green, Ky., Crawford and Associates.
- (26) Stephenson, J.B., 1989. Karst hydrology of the Kelley Spring groundwater basin with emphasis on Copas Cave: Unpublished ground-water tracing report: Bowling Green, Western Kentucky University.
- (27) Villa, R.L., 1990. Preliminary karst groundwater investigation of the area surrounding Auburn, Kentucky: Prepared for city of Auburn and Kentucky Division of Water: Bowling Green, Center for Cave and Karst Studies, Western Kentucky University, 131 p.
- (28) Aley, T., 1992. Groundwater tracing at the former Rockwell International Facility, Russellville, Kentucky: Prostem, Mo., unpublished consultant's report, 77 p.
- (29) Crawford, N.C., 1992. Investigation of karst groundwater flow and the discharge of PCBs from Armanam Spring karst groundwater basin, Simpson County, Kentucky: Bowling Green, Ky., Crawford and Associates, 14 p.
- (30) Howcroft, W.D., and Crawford, N.C., 1993. Potential sources of additional water for Auburn, Kentucky, prepared for city of Auburn and Kentucky Division of Water: Bowling Green, Center for Cave and Karst Studies, Western Kentucky University, 131 p.
- (31) Bearden, K., 1993. Dye trace from Keith Pond Spring to Lost River Rise: Unpublished ground-water tracing report: Bowling Green, Western Kentucky University.
- (32) Crawford, N.C., 1993. Unpublished ground-water tracing data: Bowling Green, Ky., Crawford and Associates.
- (33) Currens, J.C., 1993. Characterization and quantification of nonpoint-source pollutant loads in the Pleasant Grove Spring Basin, Logan County, Kentucky: A conduit-flow-dominated karst aquifer underlying an intensive-use agricultural region: Kentucky Geological Survey, Report of Investigations 1, ser. 12, 151 p.
- (34) Mull, D.S., 1993. Use of dye tracing to define the direction of ground-water flow from a superfund waste-disposal site in karst terrane, near Auburn, Kentucky: U.S. Geological Survey Water-Resources Investigation Report 92-4195, 20 p.
- (35) Crawford, N.C., 1994. Groundwater flow in the vicinity of Consumers 61 Service Station, Russellville, Kentucky: Bowling Green, Ky., Crawford and Associates, 16 p.
- (36) Crawford, N.C., and Capps, A.S., 1994. Groundwater flow in the vicinity of the Rad Chemical Company, Bowling Green, Kentucky: Unpublished ground-water tracing report: Bowling Green, Ky., Crawford and Associates.
- (37) Quinlan, J.F., 1994. Groundwater movement beneath and near Southern Sanitation, Inc., Russellville, Kentucky, as interpreted from results of tracer tests, with recommendations for groundwater monitoring: Nashville, Tenn., unpublished consultant's report, 19 p.
- (38) Paylor, R.L., 1995. Unpublished ground-water tracing data: Bowling Green, Western Kentucky University, Department of Geology and Geography.
- (39) Crawford, N.C., 1997a. Unpublished ground-water tracing data: Bowling Green, Ky., Crawford and Associates.
- (40) Crawford, N.C., 1997b. Unpublished ground-water tracing data: Bowling Green, Western Kentucky University, Department of Geography and Geology.
- (41) Crawford, N.C., 1998. Unpublished ground-water tracing data: Bowling Green, Ky., Crawford and Associates.
- (42) Crawford, N.C., 1999. Unpublished ground-water tracing data: Bowling Green, Western Kentucky University, Department of Geography and Geology.
- (43) Ray, J.A., 1999. Unpublished ground-water tracing data: Frankfort, Kentucky Division of Water.
- (44) Ray, J.A., and Nicotera, T.G., 1999. Unpublished ground-water tracing data: Frankfort, Kentucky Division of Water.
- (45) Ray, J.A., and Nicotera, T.G., 2001. Unpublished ground-water tracing data: Frankfort, Kentucky Division of Water.

**REFERENCES CITED**

AKGWA No.	Spring Name	Data Source
0050	Double	(12, 16)
0051	Cashim	(1, 3, 5, 6, 9)
0054	Lost River Rise	(2, 15, 16, 18, 31, 35, 40, 42)
0209	Kirby Mill (Hardcastle Mill)	(25)
0212	Harris	(12, 15, 39)
0286	Thomas	(9)
0310	Pleasant Grove	(33)
0315	Hobson	(33)
0360	Campbell Bluehole	(33)
0362	Clarks Bluff	(33)
0365	Stove Bluff	(33)
0369	Dunson	(33)
0369	Spring View	(33)
0489	Head of Sinking Creek	(9)
0461	Calvert	(44)
0479	Thacker	(27)
0624	Sinking Creek	(28, 35, 37)
0625	Shakertown	(30)
0629	Head of Mad River	(12)
0652	Auburn Bluehole	(24, 30)
0678	sub-basin of 0927)	(32, 41)
0705	Dorwinville	(28, 35, 37)
0925	sub-basin of 0479)	(7, 10, 11, 34)
0927	Blount	(45)
0929	Crawford	(18, 11, 27, 30, 34)
1446	Franklin	(18, 20, 23)
1447	Clear Fork Rise	(16)
1449	Jonsons Creek North	(16)
1451	Jonsons Creek South	(16)
1452	Stuenberg	(15, 14, 20)
1453	Armanam	(29)
1454	Armanam	(29)
1455	Clear Fork Spring	(20, 23)
1882	Big	(15)
1884	Kelly	(21, 26)
1885	Long Creek Road	(4, 43)
1886	Long Creek Rise	(38)
1887	Swamp Yellow	(38)
1906	Mucks	(38)



**ACKNOWLEDGMENTS:**  
We thank the many karst investigators who have contributed data for this map. Without their cooperation this map would not have been possible.

For information on obtaining copies of this map and other Kentucky Geological Survey maps and publications call:  
**Publication Sales**  
(859) 257-3896  
View the KGS World Wide Web site at [www.uky.edu/kgs](http://www.uky.edu/kgs)