

Bunting Pit B2 (Ri 31-04) May 10, 1988

Depth Elev. top Hole 17'  
(Feet)

- 0-1.5 Sand; f-m; tn; w/ brn organic-rich layers 0-0.5, 1-1.5; wood fragments common.
- 1.5-2.0 Sand; f; blk; organic  $\frac{1}{2}$  rich  
- sharp contact -
- 2.0-3.0 Sand; m-f; lt brn; sugary texture; w. sorted; stressless
- 3.0-4.5 same as above h.m. scat. common - H<sub>2</sub>O -
- 4.5-6.0 same rare h.m. lam; h.m. common
- 6.0-7.5 same poor recovery
- 7.5-8.5 same; lt gry
- 8.5-9.0 Sand; f; gry; gry cly lam common; scat. h.m.
- 9.0-10.5 Sand; f; lt gry; laminated w/ Clay; gry; some org?  
h.m. cn.
- 10.5-12.0 Clay; gry; slty; laminated w/ Sand, vf; lt gry;  
few thin organic-rich cly lam; @ 11.5-12.0 ft
- 12.0-13.5 same few sand lam; bottom 3" Sand; f; grnsh gry
- 13.5-14.0 Sand; f; grnsh grn; cly; stressless  
- sharp contact -
- 14.0-15.0 Sand; crs; grnsh gry; scat pbl (< 1cm) feld.
- 15.0-16.0 same no pbl
- 16.0-19.5 not sampled
- 19.5-22.0 Sand; crs; grnsh gry; scat h.m.; rare slily cly lam  
poor recovery
- 22.0-24.5 not sampled
- 24.5-25.0 Sand; crs; gry; scat pbl. partly wash (not sampled)
- 25.0-26.0 Sand; m grnshy grx; abn h.m.; scat h.m. lam.

- 26.0-29.5 Not sampled
- 29.5-30.0 Sand; v crs-gran; gry; <sup>common</sup> pbl ~~to~~ abn; partly wash?
- 30.0-31.0 Sand; crs; scat h.m. lam; x-bd?; gry; scat pbl; feld.
- 31.0-34.5 Not sampled
- 34.5-36.0 Same as 30-31; some Mn stains
- 36.0-39.5 Not sampled
- 39.5-41.0 same as 34.5-36.0; scat. sm l pbl.
- 41.0-44.5 Not sampled
- 44.5-46.0 Sand; crs-gran; gry; mostly wash

Augered to 55' - No change.