

GEORGIA
STATE DIVISION OF CONSERVATION
DEPARTMENT OF MINES, MINING AND GEOLOGY
GARLAND PEYTON, Director

THE GEOLOGICAL SURVEY
Bulletin Number 70

WELL LOGS OF THE
COASTAL PLAIN OF GEORGIA

by

Stephen M. Herrick, Geologist
United States Geological Survey



Prepared cooperatively by the U. S. Geological Survey

ATLANTA
1961

	Thickness (feet)	Depth (feet)
--	---------------------	-----------------

In Paleocene: Midway Group: Clayton Formation:

Limestone: gray, crystalline, fossiliferous (Foraminifera).....	100	1,040
---	-----	-------

Anomalina midwayensis at 940-950.*Operculinoides* sp. at 1010-1020.**Summary:**

No samples	450	450
In lower Eocene (Wilcox group, undifferentiated).....	260	710
No samples	230	940
In Paleocene (Clayton formation).....	100	1,040

Potential Water-Bearing Zones:

Limestone	100	1,040
-----------------	-----	-------

MITCHELL COUNTY

Location: In Sale City below elevated steel reservoir

Well No.: GGS 100

Owner: No. 1 Sale City

Elev.: 350¹

Driller: Gray Well Pump Company

Drilled: February 1940

	Thickness (feet)	Depth (feet)
--	---------------------	-----------------

No samples	315	315
------------------	-----	-----

In Upper Eocene: Jackson Group: Ocala Limestone:

Limestone: white to cream, dense, much calcitized, crystalline, fossiliferous at certain levels (Foraminifera).....	60	375
--	----	-----

Camerina striatoreticulata at 315-330.

No samples	15	390
------------------	----	-----

Limestone: as above; interbedded dolomitic limestone, dark- brown, saccharoidal	110	500
--	-----	-----

Summary:

No samples	315	315
In upper Eocene (Ocala limestone).....	185	500

Potential Water-Bearing Zones:

Limestone	185	500
-----------------	-----	-----

¹Average elevation based on Georgia State Highway Maps.