

GEORGIA
STATE DIVISION OF CONSERVATION
DEPARTMENT OF MINES, MINING AND GEOLOGY
GARLAND PEYTON, Director

THE GEOLOGICAL SURVEY
Bulletin Number 70

WELL LOGS OF THE
COASTAL PLAIN OF GEORGIA

by

Stephen M. Herrick, Geologist
United States Geological Survey



Prepared cooperatively by the U. S. Geological Survey

ATLANTA
1961

CRISP COUNTY

Location: 0.7 mi. south of Marshall near county road

Well No.: GGS 216

Owner: No. 1 H. C. Wade

Elev.: 271

Driller: H. B. Truluck

	Thickness (feet)	Depth (feet)
No samples	10	10
In Residuum:		
Clay: mottled, sandy, lignitic, and fragments of residual limestone at depth	60	70
Upper Eocene: Jackson Group: Ocala Limestone:		
Limestone: yellowish to white, fossiliferous (macroshells, echinoid and abundant bryozoan remains, and Foraminifera) ..	90	160
<i>Siphonina jacksonensis</i> at 70-110.		
<i>Operculina mariannensis</i> at 140-150.		
<i>Lepidocyclus</i> sp. abundant at 150-160.		

Summary:

No samples	10	10
In residuum	60	70
Upper Eocene (Ocala limestone)	90	160

Potential Water-Bearing Zones:

Limestone	70	160
-----------------	----	-----

CRISP COUNTY

Location: In Arabi

Well No.: GGS 225

Owner: No. 1 Marvin McKinney

Elev.: 449

Driller: H. B. Truluck

	Thickness (feet)	Depth (feet)
No samples	80	80
In Miocene (Undifferentiated):		
Clay: yellowish-green to red (mottled), sandy, arkosic	20	100
Sand: medium to coarse-grained, angular	15	115