GEORGIA STATE DIVISION OF CONSERVATION

DEPARTMENT OF MINES, MINING AND GEOLOGY GARLAND PEYTON, Director

THE GEOLOGICAL SURVEY Bulletin Number 70

WELL LOGS OF THE COASTAL PLAIN OF GEORGIA

by

Stephen M. Herrick, Geologist United States Geological Survey



Prepared cooperatively by the U. S. Geological Survey

ATLANTA 1961

Potential Water-Bearing Zones: Sand: fine to coarse-grained 20 Sand: fine to coarse-grained 10 DODGE COUNT Location: 6 mi. northwest of Eastman Well No.: GGS 20 Owner: No. 1 M. L. Sapp Elev.: 308 Driller: H. B. Truluck Drilled: February 1952 Thickness D. (feet) (f Miocene (Undifferentiated): Clay: gray to red (mottled), sandy, limonitic 50 Sand: fine to coarse-grained, cherty 30 No samples 10 In Oligocene (Undifferentiated): Limestone: white, dense, cherty, fossiliferous (bryozoan remains and some Foraminifera) 50 Asterigerina subacuta at 110-120. Rotalia mexicana var. at 120-130. Summary: Miocene (undifferentiated) 80	and the second s	9	2
Sand: fine to coarse-grained 20 Sand: fine to coarse-grained 20 Sand: fine to coarse-grained 10 Sand: fine to coarse-grained 10 Sand: fine to coarse-grained 10 DODGE COUNT Location: 6 mi. northwest of Eastman Well No.: GGS 20 Owner: No. 1 M. L. Sapp Elev.: 308 Driller: H. B. Truluck Drilled: February 1952 Miocene (Undifferentiated): Clay: gray to red (mottled), sandy, limonitic 50 Sand: fine to coarse-grained, cherty 30 No samples 10 In Oligocene (Undifferentiated): Limestone: white, dense, cherty, fossiliferous (bryozoan remains and some Foraminifera) 50 Asterigerina subacuta at 110-120. Rotalia mexicana var. at 120-130. Summary: Miocene (undifferentiated) 80	WELL LOGS OF THE COASTAL PLAIN OF GE	ORGIA	159
Sand: fine to coarse-grained 20 Sand: fine to coarse-grained 10 Sand: fine to coarse-grained 10 Sand: fine to coarse-grained 10 DODGE COUNT Location: 6 mi. northwest of Eastman Well No.: GGS 20 Owner: No. 1 M. L. Sapp Elev.: 308 Driller: H. B. Truluck Drilled: February 1952 Thickness Driftentiated: 50 Sand: fine to coarse-grained, cherty 30 No samples 10 In Oligocene (Undifferentiated): Limestone: white, dense, cherty, fossiliferous (bryozoan remains and some Foraminifera) 50 Asterigerina subacuta at 110-120. Rotalia mexicana var. at 120-130. Summary: Miocene (undifferentiated) 80	Potential Water-Bearing Zones:		
Sand: fine to coarse-grained 20 Sand: fine to coarse-grained 10 Sand: fine to coarse-grained 10 Sand: fine to coarse-grained 10 DODGE COUNT Location: 6 mi. northwest of Eastman Well No.: GGS 20 Owner: No. 1 M. L. Sapp Elev.: 308 Driller: H. B. Truluck Drilled: February 1952 Thickness (feet) (f Miocene (Undifferentiated): Clay: gray to red (mottled), sandy, limonitic 50 Sand: fine to coarse-grained, cherty 30 No samples 10 In Oligocene (Undifferentiated): Limestone: white, dense, cherty, fossiliferous (bryozoan remains and some Foraminifera) 50 Asterigerina subacuta at 110-120. Rotalia mexicana var. at 120-130. Summary: Miocene (undifferentiated) 80	Sand: fine to coarse-grained	20	60
DODGE COUNT Location: 6 mi. northwest of Eastman Owner: No. 1 M. L. Sapp Driller: H. B. Truluck Drilled: February 1952 Miocene (Undifferentiated): Clay: gray to red (mottled), sandy, limonitic			100
DODGE COUNT Location: 6 mi. northwest of Eastman Owner: No. 1 M. L. Sapp Driller: H. B. Truluck Drilled: February 1952 Thickness (feet) Miocene (Undifferentiated): Clay: gray to red (mottled), sandy, limonitic	Sand: fine to coarse-grained	10	130
Location: 6 mi. northwest of Eastman Owner: No. 1 M. L. Sapp Driller: H. B. Truluck Drilled: February 1952 Thickness Of (feet) Miocene (Undifferentiated): Clay: gray to red (mottled), sandy, limonitic 50 Sand: fine to coarse-grained, cherty 30 No samples 10 In Oligocene (Undifferentiated): Limestone: white, dense, cherty, fossiliferous (bryozoan remains and some Foraminifera) 50 Asterigerina subacuta at 110-120. Rotalia mexicana var. at 120-130. Summary: Miocene (undifferentiated) 80	Sand: fine to coarse-grained	10	170
Location: 6 mi. northwest of Eastman Owner: No. 1 M. L. Sapp Driller: H. B. Truluck Drilled: February 1952 Thickness (feet) Miocene (Undifferentiated): Clay: gray to red (mottled), sandy, limonitic 50 Sand: fine to coarse-grained, cherty 30 No samples 10 In Oligocene (Undifferentiated): Limestone: white, dense, cherty, fossiliferous (bryozoan remains and some Foraminifera) 50 Asterigerina subacuta at 110-120. Rotalia mexicana var. at 120-130. Summary: Miocene (undifferentiated) 80			
Location: 6 mi. northwest of Eastman Owner: No. 1 M. L. Sapp Driller: H. B. Truluck Drilled: February 1952 Thickness Of (feet) Miocene (Undifferentiated): Clay: gray to red (mottled), sandy, limonitic 50 Sand: fine to coarse-grained, cherty 30 No samples 10 In Oligocene (Undifferentiated): Limestone: white, dense, cherty, fossiliferous (bryozoan remains and some Foraminifera) 50 Asterigerina subacuta at 110-120. Rotalia mexicana var. at 120-130. Summary: Miocene (undifferentiated) 80	Tr.		
Location: 6 mi. northwest of Eastman Owner: No. 1 M. L. Sapp Driller: H. B. Truluck Drilled: February 1952 Thickness Of (feet) Miocene (Undifferentiated): Clay: gray to red (mottled), sandy, limonitic 50 Sand: fine to coarse-grained, cherty 30 No samples 10 In Oligocene (Undifferentiated): Limestone: white, dense, cherty, fossiliferous (bryozoan remains and some Foraminifera) 50 Asterigerina subacuta at 110-120. Rotalia mexicana var. at 120-130. Summary: Miocene (undifferentiated) 80	•		
Owner: No. 1 M. L. Sapp Driller: H. B. Truluck Drilled: February 1952 Thickness (feet) Miocene (Undifferentiated): Clay: gray to red (mottled), sandy, limonitic 50 Sand: fine to coarse-grained, cherty 30 No samples 10 In Oligocene (Undifferentiated): Limestone: white, dense, cherty, fossiliferous (bryozoan remains and some Foraminifera) 50 Asterigerina subacuta at 110-120. Rotalia mexicana var. at 120-130. Summary: Miocene (undifferentiated) 80		DODGE CO	UNTY
Owner: No. 1 M. L. Sapp Driller: H. B. Truluck Drilled: February 1952 Thickness (feet) Miocene (Undifferentiated): Clay: gray to red (mottled), sandy, limonitic 50 Sand: fine to coarse-grained, cherty 30 No samples 10 In Oligocene (Undifferentiated): Limestone: white, dense, cherty, fossiliferous (bryozoan remains and some Foraminifera) 50 Asterigerina subacuta at 110-120. Rotalia mexicana var. at 120-130. Summary: Miocene (undifferentiated) 80			
Driller: H. B. Truluck Drilled: February 1952 Thickness (feet) Miocene (Undifferentiated): Clay: gray to red (mottled), sandy, limonitic 50 Sand: fine to coarse-grained, cherty 30 No samples 10 In Oligocene (Undifferentiated): Limestone: white, dense, cherty, fossiliferous (bryozoan remains and some Foraminifera) 50 Asterigerina subacuta at 110-120. Rotalia mexicana var. at 120-130. Summary: Miocene (undifferentiated) 80			3S 269
Drilled: February 1952 Thickness (feet) Miocene (Undifferentiated): Clay: gray to red (mottled), sandy, limonitic 50 Sand: fine to coarse-grained, cherty 30 No samples 10 In Oligocene (Undifferentiated): Limestone: white, dense, cherty, fossiliferous (bryozoan remains and some Foraminifera) 50 Asterigerina subacuta at 110-120. Rotalia mexicana var. at 120-130. Summary: Miocene (undifferentiated) 80		Elev.: 308	
Miocene (Undifferentiated): Clay: gray to red (mottled), sandy, limonitic 50 Sand: fine to coarse-grained, cherty 30 No samples 10 In Oligocene (Undifferentiated): Limestone: white, dense, cherty, fossiliferous (bryozoan remains and some Foraminifera) 50 Asterigerina subacuta at 110-120. Rotalia mexicana var. at 120-130. Summary: Miocene (undifferentiated) 80	AND AND AND ARTHUR PROPERTY OF THE POPULATION OF		
Miocene (Undifferentiated): Clay: gray to red (mottled), sandy, limonitic 50 Sand: fine to coarse-grained, cherty 30 No samples 10 In Oligocene (Undifferentiated): Limestone: white, dense, cherty, fossiliferous (bryozoan remains and some Foraminifera) 50 Asterigerina subacuta at 110-120. Rotalia mexicana var. at 120-130. Summary: Miocene (undifferentiated) 80	Drilled: February 1952	<u>*</u>	
Clay: gray to red (mottled), sandy, limonitic 50 Sand: fine to coarse-grained, cherty 30 No samples 10 In Oligocene (Undifferentiated): Limestone: white, dense, cherty, fossiliferous (bryozoan remains and some Foraminifera) 50 Asterigerina subacuta at 110-120. Rotalia mexicana var. at 120-130. Summary: Miocene (undifferentiated) 80	x .		Depth (feet)
No samples			50
In Oligocene (Undifferentiated): Limestone: white, dense, cherty, fossiliferous (bryozoan remains and some Foraminifera)			_80
In Oligocene (Undifferentiated): Limestone: white, dense, cherty, fossiliferous (bryozoan remains and some Foraminifera)	No samples	10	90
Limestone: white, dense, cherty, fossiliferous (bryozoan remains and some Foraminifera) 50 Asterigerina subacuta at 110-120. Rotalia mexicana var. at 120-130. Summary: Miocene (undifferentiated) 80			•
mains and some Foraminifera) 50 Asterigerina subacuta at 110-120. Rotalia mexicana var. at 120-130. Summary: Miocene (undifferentiated) 80	In Oligocene (Undifferentiated):		
mains and some Foraminifera) 50 Asterigerina subacuta at 110-120. Rotalia mexicana var. at 120-130. Summary: Miocene (undifferentiated) 80	Limestone: white, dense, cherty, fossiliferous (bryozoan	ı re-	2.
Asterigerina subacuta at 110-120. Rotalia mexicana var. at 120-130. Summary: Miocene (undifferentiated)			140
Rotalia mexicana var. at 120-130. Summary: Miocene (undifferentiated) 80			
Summary: Miocene (undifferentiated) 80			
Miocene (undifferentiated) 80	Rotalia mexicana var. at 120-130.		
Miocene (undifferentiated) 80			
Miocene (undifferentiated) 80 No samples 10	Summary:	7	
No samples 10	Miocene (undifferentiated)	80	80
L. T.	No samples	10	90
			140

Potential Water-Bearing Zones:

30

50

Sand: fine to coarse-grained

Limestone