

GEORGIA
STATE DIVISION OF CONSERVATION
DEPARTMENT OF MINES, MINING AND GEOLOGY
GARLAND PEYTON, Director

THE GEOLOGICAL SURVEY
Bulletin Number 70

WELL LOGS OF THE
COASTAL PLAIN OF GEORGIA

by

Stephen M. Herrick, Geologist
United States Geological Survey



Prepared cooperatively by the U. S. Geological Survey

ATLANTA
1961

	Thickness (feet)	Depth (feet)
Potential Water-Bearing Zones:		
Sand: fine to coarse-grained.....	14	154
Sand: fine to coarse-grained.....	45	214
Limestone.....	21	263

Remarks:

More productive aquifers (of Upper Cretaceous age) are available below 263.

SUMTER COUNTY

Location: ½ mi. northeast of Shiloh School, at dwelling
 Owner: No. 1 Smith Moore
 Driller: Southeastern Drilling Company
 Drilled: 1952

Well No.: GGS 296
 Elev.: 509

	Thickness (feet)	Depth (feet)
--	---------------------	-----------------

Middle Eocene: Claiborne Group: Tallahatta Formation:

Sand: fine to coarse-grained, angular; interbedded clay, pale-green to tan to red (mottled), micaceous, limonitic.....	140	140
--	-----	-----

Lower Eocene: Wilcox Group (Undifferentiated):

Clay: dark-gray, silty, micaceous, abundantly glauconitic.....	14	154
--	----	-----

Summary:

Middle Eocene (Tallahatta formation).....	140	140
Lower Eocene (Wilcox group, undifferentiated).....	14	154

Potential Water-Bearing Zones:

Sand: fine to coarse-grained.....	40	140
-----------------------------------	----	-----

SUMTER COUNTY

Location: At Spring Creek, near swimming pool, at first road south of grocery
 Owner: No. 1 C. E. Pelcher
 Driller: F. P. Jones
 Drilled: May 1952

Well No.: GGS 303
 Elev.: 247

	Thickness (feet)	Depth (feet)
--	---------------------	-----------------

Residuum:

Clay: dark-brown, sandy, lignitic, and fragments of residual limestone.....	30	30
---	----	----