

INTERVAL SHEET

Page 1

VDMR Well No.: 614

Date 12/20/61

Sample Interval: from 50 to 625

PROP: Morton Frozen Foods
#3

Total depth 625

COMP: C. R. Moore

Oil Gas Water x Exploratory

COUNTY: Albemarle (Crozet)

Cuttings x Core Other

From-To	From-To	From-To	<i>Washed samples only</i>	From-To
-	-	50-	370-	-
-	-	60-	380-	-
-	-	70-	390-	-
-	-	80-	<u>400-</u>	-
-	-	90-	410-	-
-	-	100-	420-	-
-	-	110-	430-	-
-	-	120-	440-	-
-	-	Missing	450-	-
-	-	140-	460-	-
-	-	150-	470-	-
-	-	160-	480-	-
-	-	170-	490-	-
-	-	180-	<u>500-</u>	-
-	-	190-	510-	-
-	-	200-	520-	-
-	-	210-	530-	-
-	-	220-	540-	-
-	-	230-	550-	-
-	-	Missing	560-	-
-	-	250-	570-	-
-	-	Missing	580-	-
-	-	290-	590-	-
-	-	<u>300-</u>	<u>600-</u>	-
-	-	310-	Missing	-
-	-	320-	620-	-
-	-	330-	625-	-
-	-	340-	-	-
-	-	350-	-	-
-	-	360-	-	-

OWNER: Morton Frozen Foods # 3
DRILLER: C. R. Moore
COUNTY: Albermarle (Crozet)

VDMR # 614
WWCR # 450
Depth: 625'

SAMPLE EXAMINATION
(washed)

0-50	No samples
50	Weathered gneiss
50-60	"
60-70	"
70-80	" water on bedrock
80-90	Quartz-muscovite-biotite schist (70% quartz)
90-100	"
100-120	"
120-140	Biotite-muscovite schist (50% quartz)
140-150	" (60% quartz) Fracture in rock
150-160	" (45% quartz)
160-170	"
170-180	Quartz-biotite-muscovite gneiss
180-190	"
190-200	"
200-210	" with traces of epidote
210-220	Quartz-biotite-muscovite gneiss
220-230	"
230-250	"

250-290	" with some biotite schist particles
290-300	Quartz-biotite-muscovite gneiss
300-310	" with traces of epidote
310-320	Quartz-biotite-muscovite gneiss
320-330	"
330-340	"
340-350	"
350-360	"
360-370	"
370-380	"
380-400	" with traces of epidote
400-420	Quartz-biotite-muscovite gneiss
420-440	"
440-460	"
460-480	"
480-500	" with traces of epidote
500-520	Quartz-biotite-muscovite gneiss
520-540	" with traces of pyrite, small fracture in rock
540-560	Quartz-biotite-muscovite gneiss
560-580	"
580-600	"
600-620	"
620-625	"