INTERVAL SHEET

Pare 1 of 1

VDMR Well No: 1871

1

WWCR: 396

Date rec'd: 4/26/67

Sample Interval: from 0'

to: 352

PROP: E. E. Pittman

Number of samples: 26

Total Depth:

COMP: Pittman Wood & Metal Co.

Oil or Gas: Water: *Exploratory:

3521

COUNTY: Southampton (Vicksville)

Fr	om.	-To		I	Fron	n-To	Fr	om-To		From	-To
0	_	30			1.500	-		_		taria -	
30	-	92			- 1971	- 100		-		-	
92	-					_		_			
102	-				with the						
112	-					- 1		_		_	
122	-	132				_		-		-	
132		142				_		_		(i) *** _	
142		152			restle.	- 1				_	
152		162				-		_		-	
162		172			- 111			_			
		- 1 em									
172	_	182			1216	-		_ **		_	
182		192				- /		-		_	
192	-					_		_		_	
202		212				_		_		·	
212	-					_		_		_	
		the pict that									
222	_	232				_		_		-	
		mple			rel :	-		_		-	
242		252				_		_		_	
252		262				_		_			
262		272				_ " "		-		_	
402		field 2 start								1121	
272		282				_				_	
282		292				_		_		_	
292		302				_		_		_	
302		312						_		_	
312	_					_		_		_	
		top had been									
322		332				_				_	
332	_					_		_		_	
342	_	352				_		_		_	
	_					_		_		_	
	_							_		_	
									e		

^{*} Unwashed sample only

OWNER: Elmo Pittman

DRILLER: Pittman Wood & Metal Products, Inc.

COUNTY: Southampton (Vicksville)

VDMR: 1871 WWCR: 396 TOTAL DEPTH: 352'

GEOLOGIC LOG

Depth in feet

COLUMBIA GROUP (0-30')

0-30 Sand - moderately abundant matrix of bright-red clay; fine-grained, fairly well-sorted, angular; minor feldspar and muscovite

YORKTOWN FORMATION (30-142')

30-92 Clay - medium-gray, sandy; 15 percent coarse shell fragments - pelecypods, gastropods (<u>Turritella</u>), and echinoid spines; sand is fine- to very fine-grained, fairly well-sorted, angular; 3-5 percent fresh glauconite; foraminifers moderately abundant; a few bone fragments

92-102 "

102-112 Sand - dark-gray, moderately clayey; fine- to very fine-grained, well-sorted, angular to subangular; about 5 percent glauconite; minor phosphorite, and muscovite; a few shell fragments

112-122 "

122-132 "

132-142 Clay - medium- to dark-gray, sand-free to sandy; sand is fine-grained, fairly well-sorted, about 10 percent glauconite; small amounts of muscovite, phosphorite, and shell fragments

MATTAPONI FORMATION (142-172')

- 142-152 Sand abundant matrix of greenish-gray clay; very fine- to mediumgrained, moderately sorted; 65 percent quartz, 35 percent greenish-black to medium-green glauconite; phosphatic bone and shell fragments common; minor muscovite
- 152-162 Sand abundant matrix of dark clays (greens, browns, black); fineto medium-grained, fairly well-sorted; 50 percent angular
 quartz, 50 percent greenish-black to light-green glauconite;
 phosphatic nodules and bone and shell fragments common; phostrace of muscovite; a few shell fragments
- 162-172 Clay greenish-gray, micaceous, locally sandy, slightly pebbly; sand is fine- to medium-grained, fairly well-sorted; clear, angular quartz (60 percent) and dark-green glauconite (40 percent; phosphatic nodules and bone fragments common; a few shell fragments, echinoid spines, and foraminifers

TRANSITIONAL BEDS (172-202')

172-182 Sand - brownish-gray, clayey, a few small pebbles; fine- to coarse-grained, rather poorly sorted, angular to subrounded; feld-spathic; slightly glauconitic; coarse muscovite prominent; shell and bone fragments common

182-192 " fine- to medium-grained, fairly well-sorted

192-202 Sand - sparse matrix of greenish-gray, silty clay, 10 percent fine gravel; fine- to very coarse-grained, rather poorly sorted skewed fine), angular to subrounded; clear quartz, feldspathic; slightly glauconitic and micaceous; traces of garnet and phosphorite; a few shell fragments

PATUXENT FORMATION (202-352')

- 202-212 Sand light-gray, clear; medium- to coarse-grained, moderately sorted, angular to subangular; clear quartz; feldspathic; 10 percent glauconite; minor muscovite, phosphorite and garnet; a few shell fragments
- 212-222 Sand and Gravel moderately abundant matrix of brownish-gray clay;
 25 percent fine (2-8 mm), rounded gravel; 75 percent fine- to
 coarse-grained, poorly sorted sand; clear quartz; feldspathic;
 slightly glauconitic and micaceous; minor garnet; trace of shell
- 222-232 "
- 232-242 No sample
- 242-252 Sand and Gravel some brownish-gray clay; 25 percent gravel(fine);
 75 percent poorly sorted sand; clear quartz; feldspathic;
 slightly glauconitic and micaceous; minor garnet
- 252-262 Sand and Gravel fairly clean (grains are coated with white clay);
 30 percent fine (2-8 mm), rounded gravel; 70 percent coarseto very coarse-grained, fairly well-sorted, subangular to
 rounded sand; very feldspathic; trace of glauconite; minor
 garnet
- 262-272 Sand and Gravel fairly clean (grains are coated with white clay)
 30 percent fine (2-8 mm), rounded gravel; 70 percent coarseto very coarse-grained, fairly well-sorted, subangular to
 rounded sand; very feldspathic; trace of glauconite; minor
 garnet
- 272-282

282-292	Sand and Gravel - sparse matrix of white clay; 20 percent granular and very small pebbles (4-8 mm); 80 percent medium- to very coarse-grained, fairly well-sorted, subangular to subrounded sand; clear quartz; feldspathic; minor muscovite and glauconite; traces of garnet and kyanite
292-302	11
302-312	Gravel and Sand - sparse matrix of light-gray clay; 60-70 percent fine (2-20 mm) rounded gravel; 30 to 40 percent coarse- to very coarse-grained, fairly well-sorted, subrounded to rounded sand; very feldspathic; minor garnet; a few shell fragments
312-322	Clay - grayish-yellow, very sandy, slightly pebbly; sand is poorly sorted, feldspathic; minor muscovite; a few shell fragments
322-332	Sand - light-gray, slightly clayey, a few small pebbles; fine- to very coarse-grained, poorly sorted, poorly rounded; feldspathic; muscovite prominent; minor glauconite and garnet
332-342	Sand - medium-gray, slightly clayey, 25 percent granule gravel; coarse- to very coarse-grained, fairly well-sorted, angular to sub- rounded; clear quartz, feldspathic; traces of muscovite, garnet, pyrite, and glauconite; a few shell fragments
342-352	" 15 percent granule gravel

GEOLOGIC SUMMARY

	ROCK Unit	<u>Age</u>
0-30'	Columbia Group	post-Miocene
30-142'	Yorktown Formation	Miocene
142-172	Mattaponi Formation	Paleocene - Late Cretaceous
172-202'	Transitional Beds	Late Cretaceous
202-352	Patuxent Formation	Early Cretaceous

Virginia Division of Mineral Resources Robert H. Teifke, Geologist October 16, 1967

Robert H, Teifke March 3, 1972