

Generalized Geologic Map for Land-Use Planning: Lyon County, Kentucky

E. Glynn Beck, David A. Williams, and Daniel I. Carey

Acknowledgments

Thanks to John Rudolph, Lyon County Agriculture and Natural Resource agent; and James Garner, Western Kentucky Correctional Complex, for photo assistance.


EXPLANATION

- School
- Urban service boundary
- Watershed divide
- Wetlands > 1 acre, (U.S. Fish and Wildlife Service, 2003)
- Water
- Artificial fill
- Sinkholes
- Watershed divide
- Concealed faults
- Faults
- Secondary faults
- Water Walls**
 - Domestic
 - Industrial
 - Public
 - Monitoring
- 20-foot contour interval

Kuttawa Marina is one of three marinas on Barkley Lake in Lyon County. The other two marinas are the Eddy Creek Marina and the Buzzard Rock Marina. Photo by Glynn Beck, Kentucky Geological Survey.



U.S. Army Corps of Engineers bank stabilization project along the Cumberland River in Lyon and Livingston Counties. Photo by Glynn Beck, Kentucky Geological Survey.



An example of the gently rolling topography in Lyon County, which is excellent for row crop agriculture. Other parts of Lyon County may have steep slopes with narrow valleys. Photo by Glynn Beck, Kentucky Geological Survey.



Barnett Spring is a karst spring, which flows year around. Karst springs are common throughout Lyon County. Photo by Glynn Beck, Kentucky Geological Survey.



Groundwater
In the eastern and northern two-thirds of Lyon County, more than three-quarters of the drilled wells in the uplands are adequate for a domestic supply. Yields as high as 50 gallons per minute have been reported from wells penetrating large solution channels. In the low-lying areas of Eddy Creek most wells are inadequate for domestic use unless the well intercepts a major solution opening in the limestone; in that case, the yield could be very large. Wells in the uplands of the Land Between the Lakes in the southwestern quarter of the county generally do not yield enough water for domestic use. In the lowlands adjacent to Kentucky and Barkley Lakes, however, three-fourths of the wells located within the bedrock yield enough for a domestic supply and can sometimes produce more than 5 gallons per minute. Springs with flows ranging from a few gallons per minute to 1,280 gallons per minute are found in the county. Minimum flow generally occurs in early fall, maximum flows in late winter. For more information on groundwater in the county, see Carey and Stickney (2001).




PLANNING TABLE DEFINITIONS
FOUNDATION AND EXCAVATION
The terms "earth" and "rock" excavation are used in the engineering sense; earth can be excavated by hand tools, whereas rock requires heavy equipment or blasting to remove.


LIMITATIONS
Slight—A slight limitation is one that commonly requires some corrective measure but can be overcome without a great deal of difficulty or expense.
Moderate—A moderate limitation is one that can normally be overcome but the difficulty and expense are great enough that completing the project is commonly a question of feasibility.
Severe—A severe limitation is one that is difficult to overcome and commonly is not feasible because of the expense involved.

LAND USES
Septic tank disposal system—A septic tank disposal system consists of a septic tank and a filter field. The filter field is a subsurface tile system laid in such a way that effluent from the septic tank is distributed with reasonable uniformity into the natural soil.
Residences—Ratings are made for residences with and without basements because the degree of limitation is dependent upon ease and required depth of excavation. For example, excavation in limestone has greater limitation than excavation in shale for a house with a basement.
Highways and streets—Refers to paved roads in which cuts and fills are made in hilly topography, and considerable work is done preparing subgrades and bases before the surface is applied.
Access roads—These are low-cost roads, driveways, etc., usually surfaced with crushed stone or a thin layer of blacktop. A minimum of cuts and fills are made; little work is done preparing a subgrade, and generally only a thin base is used. The degree of limitation is based on year-around use and would be less severe if not used during the winter and early spring. Some types of recreation areas would not be used during these seasons.
Light industry and malls—Ratings are based on developments having structures or equivalent load limit requirements of three stories or less, and large paved areas for parking lots. Structures with greater load limit requirements would normally need footings in solid rock, and the rock would need to be core drilled to determine presence of caverns, cracks, etc.
Intensive recreation—Athletic fields, stadiums, etc.
Extensive recreation—Camp sites, picnic areas, parks, etc.
Reservoir areas—The floor of the area where the water is impounded. Ratings are based on the permeability of the rock.
Reservoir embankments—The rocks are rated on limitations for embankment material.
Underground utilities—Included in this group are sanitary sewers, storm sewers, water mains, and other pipes that require fairly deep trenches.


7.5-Minute Quadrangle Maps



Geology of Kentucky




LYON COUNTY




Copyright 2004 by the University of Kentucky, Kentucky Geological Survey
For information on obtaining copies of this map and other Kentucky Geological Survey maps and publications call:
Public Information Center
859.257.3896
877.778.7827 (toll free)
View the KGS World Wide Web site at:
www.uky.edu/kgs

Scale 1:48,000
1 inch = 3/4 mile



For Planning Use Only

This map is not intended to be used for selecting individual sites. Its purpose is to inform land-use planners, government officials, and the public in a general way about geologic bedrock conditions that affect the selection of sites for various purposes. The properties of thick soils may supercede those of the underlying bedrock and should be considered on a site-to-site basis. At any site, it is important to understand the characteristics of both the soils and the underlying rock. For further assistance, contact the Kentucky Geological Survey, Western Kentucky Office, 1401 Corporate Court, Henderson, KY 42420, phone 270.827.3414 or 270.827.3404. For more information, and to make custom maps of your local area, visit our Land-Use Planning Internet Mapping Web Site at kgsmap.uky.edu/website/kyuplan/viewer.htm.




Geologic Hazards

Many concealed and exposed faults are located throughout Lyon County. Currently, there is no evidence that these faults are active. Because of the proximity of Lyon County to the New Madrid Seismic Zone, however, strong earthquake activity is a possibility. Soil creep, slumps, and landslides along steep slopes may occur from erosion or ground motion associated with a strong earthquake.

Areas associated with saturated alluvium (unit 1) and other unconsolidated deposits (units 2 and 3) are subject to liquefaction during a strong earthquake. Alluvium deposits are also subject to flooding. Soils derived from alluvium deposits have a moderate to high shrink-swell capacity, which may affect structural foundations and roads. Flood information is available from the Kentucky Division of Water, Flood Plain Management Branch, www.water.ky.gov/floods/.

Earthquake Hazard Information

Peak ground acceleration at the top of rock that will probably occur in the next 500 years in Kentucky




Although we do not know when and where the next major earthquake will occur, we do know that an earthquake will cause damage. Severity depends on many factors, such as earthquake magnitude, the distance from the epicenter, and local geology. Information on earthquake effects is obtained by monitoring earthquakes and performing research. Such information is vital for earthquake hazard mitigation and risk reduction.

The most important information for seismic-hazard mitigation and risk reduction is ground-motion hazard. One way of predicting ground-motion hazard is by determining the peak ground acceleration (PGA) that may occur in a particular timeframe. The map above shows the PGA at the top of bedrock that will likely occur within the next 500 years in Kentucky (Street and others, 1996). It shows, as expected, that PGA would be greatest in far western Kentucky near the New Madrid Seismic Zone. Ground-motion hazard maps for the central United States and other areas are available from the U.S. Geological Survey. These maps are used to set general policies on mitigating damage. For example, maps produced by the USGS in 1996 were used to determine seismic design in building codes. For additional information pertaining to earthquake hazards, visit the Kentucky Geological Survey Web site at www.uky.edu/KGS/geologic/hazards/geologic/hazards.html.

Karst

The term "karst" refers to a landscape characterized by sinkholes, springs, sinking streams (streams that disappear underground), and underground drainage through solution-enlarged conduits or caves. Karst landscapes form when slightly acidic water from rain and snow-melt seeps through soil cover into fractured and soluble bedrock (usually limestone, dolomite, or gypsum).

Sinkholes are depressions on the land surface where water drains underground. Usually circular and often funnel-shaped, they range in size from a few feet to hundreds of feet in diameter. Springs occur when water emerges from underground to become surface water. Caves are solution-enlarged fractures or conduits that are large enough for a person to enter.



Never use sinkholes as dumps. All waste, but especially pesticides, paints, household chemicals, automobile batteries, and used motor oil, should be taken to an appropriate recycling center or landfill.

Make sure runoff from parking lots, streets, and other urban areas is routed through a detention basin and sediment trap to filter it before it flows into a sinkhole.

Make sure your home septic system is working properly and that it's not discharging sewage into a crevice or sinkhole.

Keep cattle and other livestock out of sinkholes and sinking streams. There are other methods of providing water to livestock.

See to it that sinkholes near or in crop fields are bordered with trees, shrubs, or grass "buffer strips." This will filter runoff flowing into sinkholes and also keep filled areas from sinkholes.

Construct waste-holding lagoons in karst areas carefully, to prevent the bottom of the lagoon from collapsing, which would result in a catastrophic emptying of waste into the groundwater.

If required, develop a groundwater protection plan (410KAR5.037) or an agricultural water-quality plan (KRS24.71) for your land use.

(From Currens, 2001)

Additional Planning Resources

Listed below are Web sites for several agencies and organizations that may be of assistance with land-use planning issues in Lyon County:

- www.lyoncounty.com—Lyon County
- ces.uky.edu/kyu—University of Kentucky Cooperative Extension Service
- www.peadd.org—Pennyrite Area Development District
- www.thinkkentucky.com/development/tw127—Kentucky Economic Development Information System
- www.uky.edu/KentuckyAtlas21143.html—Kentucky Atlas and Gazetteer
- quickfacts.census.gov/qfd/states/21/21143.html—U.S. Census data
- kgsweb.uky.edu/download/kgsplanning.htm—Planning information from the Kentucky Geological Survey

References Cited

- Carey, D.I., and Stickney, J.F., 2001. Groundwater resources of Lyon County, Kentucky. Kentucky Geological Survey, Open-File Report OF-01-72, 24 p.
- Currens, J.C., 2001. Protecting Kentucky's karst aquifers from nonpoint-source pollution. Kentucky Geological Survey, ser. 12, Map and Chart 27, 1 sheet.
- Hettinger, C.P., 2003. Spatial database of the Burlington Point quadrangle, western Kentucky. Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-471.
- Adapted from Fox, K.F., Jr., and Olive, W.W., 1966. Geologic map of the Birmingham Plant quadrangle, western Kentucky. U.S. Geological Survey Geologic Quadrangle Map GQ-471, scale 1:24,000.
- Hettinger, C.P., and Solis, M.P., 2003. Spatial database of the Mont quadrangle, Kentucky. Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-305.
- Adapted from Weis, P.L., and Theobald, P.K., 1964. Geology of the Mont quadrangle, Kentucky. U.S. Geological Survey Geologic Quadrangle Map GQ-305, scale 1:24,000.
- Humphrey, M.E., 1981. Soil survey of Lyon and Trigg Counties, Kentucky. U.S. Department of Agriculture, Soil Conservation Service, 147 p.
- Martin, S.L., 2002a. Spatial database of the Dycusburg quadrangle, western Kentucky. Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-1149.
- Adapted from Amos, D.H., and Hays, W.H., 1974. Geologic map of the Dycusburg quadrangle, western Kentucky. U.S. Geological Survey Geologic Quadrangle Map GQ-1149, scale 1:24,000.
- Martin, S.L., 2002b. Spatial database of the Eddyville quadrangle, Kentucky. Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-255.
- Adapted from Rogers, W.B., 1963. Geology of the Eddyville quadrangle, Kentucky. U.S. Geological Survey Geologic Quadrangle Map GQ-255, scale 1:24,000.
- Martin, S.L., 2002c. Spatial database of the Fredonia quadrangle, western Kentucky. Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-607.
- Adapted from Rogers, W.B., and Hays, W.H., 1967. Geologic map of the Fredonia quadrangle, western Kentucky. U.S. Geological Survey Geologic Quadrangle Map GQ-607, scale 1:24,000.
- Martin, S.L., 2002d. Spatial database of the Grand Rivers quadrangle, Kentucky. Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-328.
- Adapted from Hays, W.H., 1964. Geology of the Grand Rivers quadrangle, Kentucky. U.S. Department of Agriculture, Soil Conservation Service, 147 p.
- Martin, S.L., 2003. Spatial database of the Fairdealing quadrangle, Kentucky. Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-320.
- Adapted from Wolfe, E.W., 1954. Geology of the Fairdealing quadrangle, Kentucky. U.S. Geological Survey Geologic Quadrangle Map GQ-320, scale 1:24,000.
- Nelson, H.L., Jr., 2003a. Spatial database of the Lamasco quadrangle, western Kentucky. Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-618.
- Adapted from Sample, R.D., 1967. Geologic map of the Lamasco quadrangle, western Kentucky. U.S. Geological Survey Geologic Quadrangle Map GQ-618, scale 1:24,000.
- Nelson, H.L., Jr., 2003b. Spatial database of the Princeton West quadrangle, Kentucky. Kentucky Geological Survey, ser. 12, Digitally Vectorized Geologic Quadrangle Data DVGQ-385.
- Adapted from Sample, R.D., 1965. Geology of the Princeton West quadrangle, Kentucky. U.S. Geological Survey Geologic Quadrangle Map GQ-385, scale 1:24,000.
- Paylor, R.L., Flores, L., Caudill, M., and Currens, J.C., 2004. A GIS coverage of karst sinkholes in Kentucky. Kentucky Geological Survey, ser. 12, Digital Publication 5, 1 CD-ROM.
- Street, R., Wang, Z., Hank, I., and Allen, D., 1996. Source zones, recurrence rates, and time histories for earthquakes affecting Kentucky. Kentucky Transportation Center, University of Kentucky, KCT-96-4.
- U.S. Fish and Wildlife Service, 2003. National Wetlands Inventory, www.fws.gov/ [accessed 05/24/04].
- Woods, A.J., Omerick, J., Andrews, W.M., Call, S.M., Comstock, J.A., and Taylor, D.D., 2002. Ecoregions of Kentucky (color poster with map, descriptive text, summary tables, and photographs). U.S. Geological Survey, scale 1:1,000,000.